Surrey Nature Partnership
Healthy Environment | Healthy People | Healthy Economy

SURREY IMPORTANT GRASSLANDS INVENTORY

1.0 - January 2026

Surrey Nature Partnership

Surrey Wildlife Trust

SBIC
SURREY BIODIVERSITY INFORMATION CENTRE

Surrey Botanical Society

Surrey Important Grasslands Inventory 1.0

Contents:

1.	**Background**		**3**
	1.1	**Project origination & context**	**3**
	1.2	**Importance of biodiverse grassland habitats**	**3**
	1.3	**Land use**	**6**
	1.4	**Grassland criteria used for the purposes of this inventory**	**6**
2.	**Aim & objectives**		**7**
	2.1	**Overview**	**7**
	2.2	**Relevance to Surrey Nature Recovery & Biodiversity Net Gain**	**8**
3.	**Recognition methodology**		**8**
	3.1	**Using existing data-sets**	**8**
	3.2	**Ground-truthing & field surveys**	**9**
4.	**Planning policy**		**10**
	4.1	**Regulating Biodiversity Net Gain**	**10**
	4.2	**Model policy & recommendation**	**11**
5.	**Management**		**11**
	5.1	**Objective**	**11**
	5.2	**Generic management approaches**	**12**
		5.2.1 Annual bio-mass removal	**12**
		5.2.2 Applications	**12**
		5.2.3 Drainage	**13**
		5.2.4 Seeding	**13**
6.	**Summary Inventory 1.0**		**14-22**
7.	**References & further reading**		**22**

Annex 1. Surrey golf courses with significant grassland biodiversity interest
Annex 2a. Grassland axiophytes in Surrey used in methodology
Annex 2b. CHEGD fungi species in UK
Annex 3. Waiting List of potential, as yet unverified sites of future interest

[SEPARATE DOCUMENT]
Appendix I. Inventory 1.0, inc. maps
Appendix II. Inventory 1.0; key statistics

Cover photo: Surrey's Coronation Meadow at The Sheepleas SSSI

1. Background

1.1 Project origination & context

The Surrey Important Grassland Inventory is the first project of its kind to document species-rich grasslands across a whole county. For over a decade there has been a collective desire within Surrey's nature conservation sector to comprehensively document the remaining biodiverse grasslands across the administrative county. The **Surrey Botanical Society** (SBS) was a strong proponent in this. The project was enabled when part-funding became available in 2022 from **Natural England**'s Nature Recovery Network[1], which was matched in kind by **Surrey Wildlife Trust** (SWT) with contributions from **Surrey Biodiversity Information Centre** (SBIC). In 2022-3 additional funding was awarded by both the **Banister Charitable** and the **Hamamelis** Trusts, specifically to extend the project's programme of original survey and re-survey (see 3.2). A project steering group was formed comprised of relevant representatives from the **Surrey Nature Partnership** (SyNP), which would publish the product.

1.2 Importance of biodiverse grassland habitats

The Botanical Society of Britain & Ireland (BSBI) is the longest-standing champion for the research and conservation of Britain's flora. Findings from the culmination of their *Plant Atlas 2000* project are summarised in the document *Britain's Changing Flora: A Summary of the Results of Plant Atlas 2020* (BSBI 2023)[2]. This finds that: "*..our native flora is much diminished relative to the situation recorded by our predecessors in the 1950s*." And "*..a holistic plan of action to reverse this decline, ..should focus on strengthening the protection of species-rich areas, ..and reducing damaging land use practices*." According to the document, "*Grasslands are amongst the most threatened habitats in Britain, mainly because of changes in agriculture since the 1950s*" (see **Figure 1**).

Internationally, there has been growing pressure from major environmental non-governmental organisations (eNGOs) for their governments to value grasslands as critical ecosystems for nature, climate and people[3]; to be realised through coordinated action to develop integrated policies for protection and sustained conservation of those grasslands of highest ecological importance. In the UK the eNGO Plantlife, a champion of the value of grasslands for their biodiversity and ecosystem services, published a 2024 discussion paper on ancient grasslands by the British Ecological Society (BES) to create a practical definition to better identify and protect important grassland types[4].

Grasslands support many invertebrates responsible for **crop and fruit pollination**, and for the maintenance of healthy soils. The significance of grasslands to **carbon sequestration** is gaining better recognition, with the oldest and richest examples exceeding that of many woodlands. Grasslands on floodplains are becoming increasingly essential for the absorption of seasonal peaks in river water-levels, **buffering potential flooding events** both

[1] The Nature Recovery Network - GOV.UK

[2] See; Walker, K.J., Stroh, P.A., Roy, D.B., Burkmar, R.J. & Prescott, O.L. (2003) *Britain's Changing Flora. A Summary of the Results of Plant Atlas 2020.* BSBI

[3] Grasslands, Rangelands, Savannahs & Shrublands Alliance (2024). Discussion paper on Valuing Grasslands: Critical Ecosystems for nature, climate and people.

[4] British Ecological Society (2024). Discussion Paper on Ancient Grasslands. Plantlife.

within the immediate environment and downstream.

There has been a definitive national inventory of 'ancient' woodlands[5] since the early 1980s administered by Natural England (NE) that informs national and local planning policy protecting ancient woodland in development management. Ancient woodlands are defined as currently wooded sites with evidence to support their presence from CE 1600 to the present day, the importance of which is inferred from the antiquity and undisturbed quality of their underlying soils and the assumed continuity of a woodland canopy. Similar to the ancient woodland inventory providing for our valuable woodlands, this Surrey grasslands inventory aims to provide a practical resource to raise awareness of our most valuable grasslands, to elevate their bespoke protection through the planning system in a similar way.

The grasslands selected for this inventory could be described variously as ancient/old, historic/heritage, un- or semi-improved, species-rich and/or biodiverse. We have chosen to adopt 'Important', as a generic qualifier that can be widely understood by specialists and non-specialists alike. The inventory also recognises the hidden value of essentially undisturbed, below-ground biodiversity within the different habitat classifications based on above-ground vascular plant diversity. By compiling this important grassland inventory we provide a tool to emphasise their value as complex ecosystems, that provide a fundamental role throughout our landscapes to ensure planetary and human health, while also of related importance in evaluations of natural capital.

Figure 1. Showing the smoothed trends (1930-2019) for plant species typically associated with acid, calcareous and neutral grasslands in Britain.
[reproduced by kind permission of the BSBI]



[5] Revised Ancient Woodland Inventory for Surrey (2011) – in A review of the revision of the Ancient Woodland Inventory in the South East (NERR042), Natural England 2011.

Broadleaved woodland is estimated to occupy at least 21% of Surrey but in contrast, un- or semi-improved species-rich grassland occupies a much lower percentage. Surrey's least agriculturally improved permanent grasslands support a unique and declining biodiversity, including plants, a great many invertebrates and some of their ecologically dependent vertebrates. It is the long-term undisturbed soils of these least improved grasslands that are responsible for supporting diverse assemblages of fungi, belonging especially to the 'CHEGD' families and genera[6].

Loss of biodiverse grasslands has occurred over time whether through disturbance, degradation or through cyclical or permanent conversion to arable, as generic agricultural 'improvement' driven by its primary purpose: to grow more food, more quickly. Many have had their soils altered through elevations in fertility and drainage. In the past, if too acidic their pH was frequently elevated by liming, while fertility has been continuously supplemented and replenished using manures, treated sewage sludge-cake, synthetic N:P:K inputs or liquid digestates. Pasture has been repeatedly re-seeded with high productivity, protein-rich grass cultivars; and herb- and pesticides are routinely applied to eliminate the remaining pernicious 'weeds', fungi, and herbivorous beetles, bugs and molluscs.

Where corners of our farmed landscapes have escaped this relentless improvement impacting their biodiversity down the centuries, those fields used longest as grazed pastures or mown meadows where this impact has had least effect are termed 'semi-natural' grasslands for biodiversity purposes. These are then further classified by their soil pH range (see **Box 1**). For various reasons - both historic and environmental combined, Surrey has suffered less than many comparable lowland counties in this regard, certainly within the southern half of the county.

Box 1

Acid grasslands (both dry and ephemerally wet) can occur as enclosed fields or on commons and village greens as the predominant vegetation type over sand and gravelly substrates both north and south, and on top of the chalk escarpment. They are excluded from this inventory where they occur within the complex mosaic of intergrading habitat-types on heathlands, however.

Calcareous grasslands are principally found over the Chalk of the North Downs, but also within the immediate Thames floodplain where calcareous river-borne silt carried from higher in the catchment has had an influence on their pH and composition. Potentially, both the Bargate series of the Wealden Greensands and the *Paludina* limestone of the Low Weald could also support calcareous grassland, but most exposures of these rocks are wooded in Surrey.

Neutral grasslands generally occur over clay substrates mainly in the Low Weald and on river floodplains. These grasslands have almost disappeared from modern lowland landscapes as a result of agricultural improvement and only a few very rare examples now remain in Surrey as pasture and/or traditional 'hay meadows'.

[6] CHEGD = **C**lavarioids, ***H**ygrocybe* (waxcap species), ***E**ntoloma*, ***G**eoglossum* & allies (the earthtongues) & ***D**ermoloma* species (see Annex 2b, and Plantlife's guide to waxcap fungi).

1.3 Land use

A significant proportion of our formerly farmed landscapes is now used for recreation. This includes widespread equestrianism and conversion to golf courses. Indeed, grassland within golf courses can serve to preserve species-richness; in this inventory these are included in a separate category in Annex 1 rather than in the main inventory[7], as these areas are often managed as roughs or fairways with little fixed delineation.

Historic parkland can contain remnant undisturbed old grassland as can commons and village greens sometimes with cricket pitches, where any former agricultural function has for the most part been superseded. Despite being presently combined with the habitat type 'wood-pasture' and thus broadly defined as woodland for purposes of habitat classification, un- and semi-improved grassland in historic parklands is included in this inventory.

The long-surviving and undisturbed species-rich grasslands enclosed within some ancient churchyards and cemeteries have also been included in this inventory and can be very important for grassland fungi, as well as for a range of other wildlife species[8]. The efforts of the Eco Churches (A Rocha UK) and Caring for God's Acre initiatives have fostered an awareness and conservation of this interest.

1.4 Grassland criteria used for the purposes of this inventory

1. The Surrey Important Grasslands Inventory (henceforth the Inventory) includes eligible grassland land parcels that can be provisionally classified as representative of a selected series of qualifying habitat categories[9] (see **Figure 2**), based on survey information currently available.

2. The qualifying habitat category may not be entirely present across the parcel but must be extensive, although there is no minimum site size/area threshold limiting inclusion on this inventory[10].

3. Where present as minor components within a complex mosaic of other non-qualifying habitat categories, which furthermore lack a defined boundary, these otherwise qualifying grasslands are **excluded** from the Inventory. Examples include a typical situation of acid grassland patches amongst lowland heathland, and grassland within open woodland glades and rides. This is also the criterion that sub-categorises qualifying habitat within golf courses, as explained in 1.3 above.

4. Species-rich grassland occurring within the margins of transport infrastructure is also **excluded** from the Inventory (but see note on their inclusion within 'Conservation Verges'[11]).

[7] Note: this does not imply that such habitats surviving in golf courses are of insufficient quality (see list in Annex 1); as with acid grassland on heathlands (see 1.5), the issue primarily concerns their geographic imprecision.
[8] See; Eco Churches and Caring for God's Acre.
[9] See UK Habitat Classification 2.0 (pre-registration required) for individual habitat identification criteria, including type codes g1a/d; g2a/c; g3a; & selectively g3c. It should be noted that these criteria include the attainment of threshold counts of certain listed Indicator Species, not all of which occur in Surrey or are very rare.
[10] This is because i) Surrey's land parcel size is significantly below the national average; and ii) there is a strong correlation of smaller fields retaining richer plant communities in most parts of the county.
[11] Grasslands supporting species of conservation concern on roadside verges have been identified separately for a 'Conservation (Road) Verge Habitats Action Pan' (SCC 2025), with a key objective to actively manage these as Roadside Nature Reserves in the future.

2. Aim & objectives

2.1 Overview

The **aim** of the Surrey Important Grassland Inventory is to increase awareness of the location and importance of Surrey's remaining species-rich grasslands, putting these on a par with ancient woodland; thereby promoting their protection in the planning system, as well as their maintenance, restoration and enhancement in land management; by using the opportunities realisable as off-site net gain projects, and in strategic natural capital auditing; within the modern administrative county of Surrey.

Figure 2. Grassland habitats qualifying for the Important Grassland Inventory, based on UK Habitat Classification 2.0[9]

UK Habitat Classification 2.0				
Level 2	**Level 3**	**Level 4**	**Level 5**	**'Distinctiveness' used to evaluate Biodiversity Net Gain (BNG)**
		(inc. affinities with S.41/Priority habitats)	**(*=European designated habitats)**	
Grassland	Acid grassland	Lowland dry acid grassland **g1a** = Priority habitat	Other lowland dry acid grassland	very high
		Other lowland acid grassland **g1d**		medium
	Calcareous grassland	Lowland calcareous grassland **g2a** = Priority habitat	***Dry grasslands & scrub on chalk or limestone; lowland**	high
			***Dry grasslands & scrub on chalk or limestone; important orchid sites**	
		Other calcareous grassland **g2c**		(*not currently included in Metric*)
	Neutral grassland	Lowland meadows **g3a** = Priority habitat	Lowland hay meadows	very high
		Other neutral grassland **g3c** [in BNG 'Good' Condition only] (inc. **Floodplain grazing-marsh** = Priority habitat)	*Arrhenatherum* neutral grassland	medium
			Lolium-Cynosurus neutral grassland	
			Deschampsia neutral grassland	
			Holcus-Juncus neutral grassland	

This aim is achieved through the following **objectives**:

- Establishing a **Surrey Grassland Steering Committee**[12] to direct the inaugural publication, and subsequently to review and advise on revisions (see last).
- Presentation of the Inventory as an evidence base, to support bespoke **protective planning policy** recommended by the Surrey Nature Partnership for adoption by Surrey's planning authorities (see 4.2).
- Supporting land managers by sharing knowledge of the importance of their species-rich grasslands, as well as best practice for their maintenance; and encouraging grassland restoration and creation projects.

[12] Representation to include Surrey Wildlife Trust, Surrey Biodiversity Information Centre, Surrey County Council, Natural England, Surrey Botanical Society, Surrey Butterfly Conservation and Surrey Fungus Study Group (all members of the Surrey Nature Partnership).

- Data-sharing of the Inventory to align with all respective and corresponding datasets, including the Surrey Local Nature Recovery Strategy (see 2.2) and the Natural England **Priority Habitat Inventory**[13] as presented on the MAGIC.gov website, where possible.
- Designation of hitherto unrecognised species-rich grasslands as **Sites of Nature Conservation Importance** (SNCI), for their grassland habitat interest. The Inventory will also serve to re-confirm existing SNCI for their grassland habitat interest.
- Subsequent revisions of the Inventory (at <10 year intervals), to include the addition and/or removal of sites as and when new data may be collected, for example through enhanced site access.

2.2 Relevance to Biodiversity Net Gain & Surrey Nature Recovery

Figure 2 presents the qualifying habitat categories in the Inventory, also showing how they relate to Priority habitats[14]. The table also refers to the ranked recognition of the relative 'Distinctiveness' of these grassland habitats within the current 'Statutory' version of the Defra Metric, used to evaluate Biodiversity Net Gain (BNG). From this it may be seen that only 'high' and 'very high' distinction types are included, plus two 'medium' types if/when their 'Condition' is rated to be 'good'. The potential relevance and use of this detail is explained in 4.1 below.

The Environment Act 2021 introduced a legal requirement to produce Local Nature Recovery Strategies (LNRS) and in Surrey the appointed *responsible authority* for this was Surrey County Council. LNRS are required to map the current distribution of various habitat types as components of local 'nature recovery networks'. In addition to datasets such as the Revised Ancient Woodland Inventory, this Inventory is being used by SCC to inform their mapping requirement.

3. Recognition methodology

3.1 Use of existing datasets

Previous county-wide surveys have identified grassland habitats (see **Figure 3**). The first stage of the Inventory used these datasets, where relevant, to help inform compilation of an initial draft list of species-rich grassland sites. A subsequent synthesis selected qualifying sites using a process including:

1. Use of the **Surrey Botanical Society** (SBS) **database** to extract records of a recognised set of grassland **axiophyte**[15] species typifying 'good quality' grassland in Surrey (axiophytes are listed in Annex 2a).
2. Creation of a raw map showing incidence point data for concentrations of both 10-19,

[13] See; https://environment.data.gov.uk/dataset/92b43165-0dd0-4e69-a712-1e49bb5aa0d0

[14] Also known as Habitats [and species] of principal importance in England - GOV.UK

[15] **Axiophytes** are species that are strongly indicative for determining conservation priorities. They should not be confused with the concept of rarity or conservation status. They may also be rare or scarce, but their signal of habitat *condition* is of importance here. Sites with more axiophytes are usually of greater importance than those with less; and monitoring changes in the diversity/extent of axiophytes on a site can indicate the success (or otherwise) of its management (Axiophytes - BSBI).

and ≥20, historical axiophyte records across Surrey; followed by a geographic analysis of results to ascertain the actual land parcels hosting these records.

3. The land parcels identified were then reviewed to assess whether records could be accurately ascribed to them or, for example, from peripheral transport infrastructure verges. These, plus sites arising in other excluded land-uses (see 1.3) were then removed.
4. The resulting list of land parcels was then cross-referenced with the initial draft site-list, to consolidate any duplications or anomalies in the site nomenclature. This became the emergent 'master', provisional Inventory.
5. A Surrey Important Grassland Inventory **GIS mapping layer** was then developed, using digital GIS shapefiles of the land parcels in the emergent Inventory (informed also by original maps from historical surveys and other datasets included in Figure 3).
6. Incorporation of additional sites into the emergent Inventory supporting important grassland fungi, as recommended by the Surrey Fungus Study Group's analyses of existing data and using expert knowledge.
7. Interpretation of **aerial imagery** (using Google Maps) to indicate the current state of sites on the emergent Inventory. This added further additional sites, albeit requiring further verification, that appeared at least as suitable as those already identified.

Figure 3. List of existing datasets

Dataset	Source	availability to the Inventory project
SNCI (re-)survey datasets, cumulative	SBIC	open access
SBS records database	SBS/SBIC	open access
BC Surrey Branch records database	Butterfly Conservation Surrey	open access
Surrey Fungus Study Group data-base/'Surreytrack'	SFSG	open access
RBGKew database (aka 'Herbtrack')	RGBK	open access
NCC Chalk Grassland Survey 1987-8	NE (as NCC)/SBIC	open access
English Nature Grassland Inventory: Surrey (1995)	NE (as English Nature)	open access
Chalk Grassland Survey, Humphreys, S. (1999)	SBIC	open access
Preliminary Botanical Survey of Unimproved Neutral & Acidic Grassland, G. Steven 1990	NE (as NCC)/SBIC	open access
ND01-03 & WG11 HPI Project Site Registers	SWT	restricted
Priority Habitat Inventory (PHI)	NE	open access
SCC/LandApp 'Best Guess' land-use dataset	SCC	open access
Survey of Old Meadow Sites in Surrey, 1979	NE (as NCC)/SBIC	open access
[*Space4Nature* – emerging]	multi-agency	restricted

3.2 Ground-truthing & field surveys

The resultant 'master' emergent Inventory was then used to develop a **survey programme**. Listed sites were prioritised for survey based on the following considerations:

- Sites with no records in the SBS database (mainly those added as appearing visually suitable from aerial photography) had highest priority for an original Inventory-initiated survey.

- Sites for which their axiophyte records were now regarded as out-of-date (made <2000) were next prioritised for re-survey.
- SNCI surveyed before 2010 were next prioritised, especially if aerial photography suggested that significant deterioration may have occurred. Conversely, SNCI or part-SNCI with recent (≥2010) survey data were regarded as not requiring re-survey.
- Sites that are/part of designated Sites of Special Scientific Interest (SSSI) were assumed to still support the qualifying habitat-type they were notified for, and not requiring re-survey. These are automatically included on the Inventory.

Sites thus prioritised made up an Inventory 'Waiting List' requiring further survey data. The Inventory survey programme commenced in 2023, when some 75 sites were surveyed. In 2025 a majority of the remaining sites from the Waiting List were surveyed. A small proportion could not be successfully accessed. The total survey effort combined professional officer time provided by SWT with voluntary time provided by members of the SBS. A bespoke Inventory App was developed by SWT to assist field surveyors in capturing the essential information.

The government Land Registry database was selectively interrogated for ownership information, to facilitate mailed requests for permission to access some private land parcels. Local knowledge has enabled further access permission to be granted, while considerable data has been collected from public Rights-of-Way passing through land parcels.

This resultant Inventory 1.0 is based on the best-available survey information available at the present time. The qualifying habitats are described as *provisional* in recognition of the variation across sites in the relative strength of the data required to validate these. We acknowledge the likelihood of there remaining further sites supporting the qualifying grassland habitats not included in this version of the Inventory, which is the primary reason for viewing it as an inaugural baseline from which future revisions may be completed to a suitable temporal cycle (see 2.1, final objective, above).

4. Planning policy

4.1 Regulating Biodiversity Net Gain

All planning permissions must now include a condition to achieve a minimum 10% net gain in Biodiversity/Hedgerow/Watercourse Units, calculated using the Statutory Metric. In the absence of a comprehensive habitat mosaic dataset for LPAs jurisdictive areas, the regulation of Statutory Metrics supporting planning applications must essentially accept as accurate the baseline evaluation claimed by their professional advisors. The Inventory will now contribute to a desirable pre-ascertained habitat mosaic dataset, to enable a more objectively informed planning regulatory system.

The Statutory Metric also incorporates an inviolable mechanism for protecting defined 'irreplaceable' habitats. While none of the grassland habitat-types of the Inventory are currently defined as irreplaceable (not even those of 'very high' distinction), subsequent revisions to BNG may in time add to this definition of irreplaceability. Meanwhile, the Surrey Nature Partnership has issued bespoke guidance for the consideration of relative

irreplaceability of Surrey's important habitats[16].

4.2 Model policy & recommendation

A model policy to feature within relevant chapters of local plans could read as follows:

> *"…Any development resulting in the loss or deterioration of biodiverse grasslands listed on the Surrey Important Grasslands Inventory (SyNP 2026, or subsequently revised) should be refused, unless there are wholly exceptional reasons and a suitable compensation strategy exists."*

Surrey's current or future LPAs are recommended to adopt such a policy within relevant chapters of their Local Plans, and other planning guidance documents, that will provide an additional constraint to development proposals impacting the sites/field parcels included on the Inventory. The material consideration thus implied, can be added to all other(s) that may also apply (such as designation as a SSSI/SNCI; presence within the Surrey Hills/High Weald National Landscapes; Green Belt; or any agricultural importance designations). Regulation of the management of sites included in the Inventory is largely outside the scope of this project, unless of course sites are also SSSI; are registered as off-site Biodiversity Net Gain enhancement sites; or to a more limited extent, are adopted as SNCI.

To an extent the Environmental Impact Assessment (EIA) (Agriculture) Regulations 2014 guard against agricultural management that could impact the biodiversity and bio-abundance of species-rich grasslands[17]. In certain circumstances, Natural England must decide if planned changes to the use of land requires a 'EIA screening decision', to determine whether these can take place. If increasing the productivity of uncultivated land (defined as untilled/fertilised for >15 years), or semi-natural land (ie. supporting Priority habitat), an EIA screening decision is necessary.

5. Management

5.1 Objective

The primary objective for management of sites listed in the Inventory is to at least maintain their botanical diversity and the abundance of this diversity. Both these values are essentially dynamic, so this maintenance is not absolute but should remain within perceptions of constancy.

Probably the majority of Inventory sites could be enhanced to some degree, however. This could be in either/both diversity and abundance values. The botanical interest of sites may be restricted to part of the field(s), see 1.4(2) above, where this might be extended to their entirety. These features influence these habitats' Statutory Metric 'condition'. Others include the presence and extent of perceived native and recognised non-native invasive species; variety in sward heights; and to an acceptable degree, the presence of bare soil

[16] See; https://surreynaturepartnership.org/wp-content/uploads/2020/08/irreplaceable-habitats-guidance-for-surrey_final_aug2020.pdf

[17] See; Environmental Impact Assessment (Agriculture) Regulations 2014.

areas (but not from excessive poaching by livestock). Just by re-introducing a suitable grazing regime, several of these factors can be improved within a single field season, thus raising the condition of the habitat.

5.2 Generic management approaches

5.2.1 Annual bio-mass removal.

Additional annual biomass removal is essential and is achieved either by mowing (with collection), grazing or a combination of both. The frequency of cuts, and/or the density of stocking are clearly adjustable and the ideal numbers to maintain/enhance the habitat is best judged from experience, although guidance literature is available[18].

As a management tool, grazing usually produces the best results in addressing most of the condition factors mentioned above. However, there are many sites where grazing is restricted as an option, usually where the level of public access (including for dog-walking) is preventative. Other difficulties may include poor or no fencing, and with sourcing on-site drinking water. Sometimes, a compromise involving temporary grazing may become possible, using portable electric fencing and bowsers. Public access would be suspended while this is in operation, but would resume alongside the former mowing regime in the future. This would likely be at relatively high grazing pressure as 'pulse grazing', used as a solution to a particular condition problem such as rank grasses or scrub removal.

Mowing should always have its arisings collected and removed from the habitat. These can be piled in a suitable location of low importance nearby, but never 'flown' so as to remain *in situ*. Impactful nutrients will accumulate if this happens, which is undesirable in all modern biodiverse grasslands. When these habitats were more widespread in the landscape, the following recommendation would have been unnecessary and probably unthinkable if functional haymaking was the objective: if a field is managed by mowing primarily as a biodiversity asset, it is advisable to leave up to 20% as a rotated uncut patch in every year, mainly to support over-wintering invertebrates. Particularly rich botanical communities, especially those of calcareous grasslands, are ideally rested from mowing or grazing at the peak of summer (at least in June) to allow an adequate period for flowering, pollination and fruiting of plants. Plantlife's recent 'No-Mow May' initiative is an attempt to raise national awareness of the need for grassland managers to rest mowing regimes at a key point in the wildflower flowering calendar.

5.2.2 Applications.

Clearly, there is no cause or reason for applying additional fertilisers over the inevitable input by any grazing stock. Also, the only herbicides to be used should be for targeted control of infestation by non-desirable invasive species and then subject to strict legal and any corporate policy constraints. The health and welfare of stock would also be managed carefully to avoid, for example, use of ivermectins due to their detrimental impact on invertebrates.

[18] See; Lowland Grassland Management Handbook (Second edition). Natural England 1999

5.2.3 Drainage.

This is only likely to apply to sites hosting g3c habitat managed as Floodplain grazing-marsh. These have been selected for inclusion in the Inventory only where whatever historic drainage regime was used in the past now appears to be largely failing, and they have become neglected as functioning grazed pasture for this or another reason. This benign neglect has allowed some recovery of their botanical diversity, sufficient to be classified as 'Other neutral grassland' and within an, at least localised, state interesting enough to merit 'good' condition.

Here, the flooding regime might be restored for them to function more as 'water-meadows', by reconnection of any former meanders or ox-bows, and the adjustment of any sluices and other remaining drainage infrastructure; in order to hold more water, more regularly and for longer periods of the year. Grazing could still take place in the summer. This form of management could be combined with any wider Natural Flood Management projects within the vicinity, that may be designed to also re-wet adjacent wet heathland or wet woodland habitats within its scope.

5.2.4 Seeding.

A management regime employing all the above recommendations would in time be expected to promote botanical diversity, abundance and if required also extent, in the mid- to long-term. There are however, techniques to accelerate this process if this is considered either desirable and/or is otherwise necessitated.

First, the seeding of native-sourced Yellow-rattle *Rhinanthus minor* into post-mown and/or lightly scarified swards will lessen the vigour of their grass component, thus promoting the germination of more abundant, as well as colonisation of additional forb species. This is usually attempted in habitats so deficient in this regard that they are unlikely to be included in this Inventory, however.

On sites where the botanical interest is confined to certain areas of a field, once the reason(s) for this community disparity is diagnosed and addressed, the mown arisings from the more diverse sections can be collected as 'green hay' (in late June-early July) and spread onto the more depauperate sections. By extension, this can be used as a technique for the restoration of adjacent or close-by, non-Inventory field parcels if the desire to do so is present.

For sites included in this Inventory, it is very unlikely that use of commercial seed-mixes, or the planting of pre-grown plants, would ever become necessary for the enhancement of their qualifying habitats. Although the availability of products marketed as being of 'local' provenance is increasing, with the implication that their use is suitable on sites of conservation importance; local provenance in this case generally refers to the national context and even their use is therefore inadvisable. Seeding or planting is therefore to be avoided where native populations of plants still thrive.

6. Summary Inventory 1.0

The table below is a summary containing the essential attributes for all sites included in the Inventory 1.0. This information is more fully presented as a separate document, with maps, in the Appendix.

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
Elmbridge (East Surrey Council >2027) = 11 sites						
TQds6705	Byfleet, Brooklands community park grasslands	TQ06	TQ06676205	g1a/d, g3c	SNCI	LA
TQrk3155	Cobham Park	TQ15	TQ11315955	g3c/fg-m	SNCI	private
TQon3600	Cobham, field W of Old Common	TQ16	TQ10366100	g1a/d	SNCI	LA
TQon0290	Cobham, Old Common	1Q16	TQ11026090	g3c	SNCI	LA
TQds5843	Cobham, The Fairmile fields	TQ16	TQ11586143	g1a/d		private
TQor4194	Hurst Park, inc. Hurst Meadows & Hurst Minor	TQ16	TQ14416894	ga/d	SNCI	LA
TQth1812	Molesey Heath	TQ16	TQ13186712	g3c	SNCI	LA
TQ E5936	West End, Garson Farm, Longmore Meadow E	TQ16	TQ12596436	g3c/fg-m		LA
TQry2250	West Molesey cemetery	TQ16	TQ13226850	g1a/d*		CoE
TQry3059	Weybridge, St.James' churchyard & cemetery	TQ06	TQ07306459	g1a/d		CoE
TQen4167	Whiteley Village, green	TQ06	TQ09416267	g1a/d*	SNCI	private
Epsom & Ewell (East Surrey Council >2027) = 9 sites						
TQrk1067	Cheam, Nonsuch Park	TQ26	TQ23106367	g3c	SNCI	LA
TQns8112	Epsom & Walton Downs	TQ25	TQ21815812	g2a/c	SNCI	private/LA
TQry5114	Epsom cemetery	TQ25	TQ21515914	g2c*	SNCI	CoE/LA
TQir7282	Epsom, covered reservoir	TQ25	TQ21725882	g2a/c	SNCI	SSEW
TQrk5965	Epsom, Livingstone Park	TQ16	TQ19596165	g3c	SNCI	LA
TQll9135	Epsom, Walton Downs-Juniper Hill	TQ25	TQ21915735	g2a/c	SNCI	LA
TQNR8992	Ewell, Howell Hill NR	TQ26	TQ23896192	g2a	SNCI	SWT
TQNR9545	Ewell, Priest Hill NR	TQ26	TQ22956145	g2c	SNCI	SWT
TQrm3497	Ewell, Warren Farm	TQ26	TQ23346297	g3c	SNCI	WT/LA
Guildford (West Surrey Council >2027) = 93 sites						
TQns3607	Albury Downs	TQ04	TQ04364907	g2a/c	SNCI	private
TQe)7432	Albury Park (composite)	TQ04	TQ06744732	g1a/d	SNCI	private
TQrd3285	Albury Park, old churchyard	TQ04	TQ06324785	g1a/d		CCT
TQrd0362	Albury, churchyard	TQ04	TQ05034762	g1a/d		CoE
TQld5995	Albury, Lacey's Field	TQ04	TQ05584695	g1a	SNCI	private
TQrm9669	Albury, meadows W of Ford Farm	TQ04	TQ04964669	g3c/fg-m	part-SNCI	private
TQld6558	Albury, Woodside Farm field	TQ04	TQ05654658	g1a		private
SUs 3453	Ash Green, Cardinals Fields	SU84	SU89344953	g3c damp	SNCI	Land Trust
SUge1267	Ash Ranges, Henley Park Range	SU95	SU93125367	g1a	SSSI	MoD
SUW)4370	Ash, Shawfield Lane meadows (W)	SU85	SU88435070	g3c	SNCI	BVCP/LA
SUge0612	Bisley, Century Range	SU94	SU94065812	g1a/d	SSSI	NRA
SUge2261	Bisley, Short Siberia Range	SU95	SU94225861	g1a	SSSI	NRA
TQn 6240	Bookham, Bank's Common	TQ15	TQ11625640	g3c damp		NT
TQrm5673	Burpham Court Farm	TQ05	TQ00565273	g3c/fg-m	SNCI	LA
TQrd8831	Chilworth, St. Martha's Hill & churchyard	TQ04	TQ02884831	g1a	SNCI	SCC, CoE
SUrd6538	Compton, Watts Mortuary Chapel churchyard	SU94	SU95654738	g1a/d		CoE
TQns6032	East Clandon, Clandon Downs	TQ05	TQ05605032	g2c	SNCI	private
TQrm2511	East Clandon, fields E of Wellhouse Farm	TQ05	TQ05255011	g2c		private
TQrk8520	East Clandon, Hatchlands Park	TQ05	TQ06855220	g3c	SNCI	NT
TQs 4688	East Horsley, Chalk Lane, field W of Greystones	TQ05	TQ09465188	g2a		private
TQse4864	East Horsley, field at Lark Rise	TQ05	TQ09485164	g2c		SCC/SWT
TQds3798	East Horsley, Rowbarns Manor fields	TQ15	TQ10375198	g2c		private
TQon3224	Effingham Common	TQ15	TQ10325524	g3c		LA
TQNR4986	Gomshall, Colekitchen Down NR	TQ04	TQ08494886	g2a	SNCI	SWT
TQ1)1064	Gomshall, Colekitchen Farm (SSSI unit 001)	TQ04	TQ09104864	g2a	SSSI	private
TQwn2238	Gomshall, field S of Hackhurst Down	TQ04	TQ09224838	g2c		private
TQds9159	Gomshall, Netley House grounds	TQ04	TQ07914859	g2a/c		NT
TQrk0720	Gomshall, Netley Park	TQ04	TQ08074820	g2c	SNCI	NT
SUse7302	Guildford, field S of Manor Copse	SU94	SU96734902	g2c		private
SUws2311	Guildford, Parsonage Meadows	SU95	SU99235111	g3c/fg-m		LA
TQwn8191	Guildford, Pewley Down	TQ04	TQ00814891	g2a/c	SNCI	LA, private

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQnd2720	Guildford, Pewley Meadows & Warren Farm/Rosamund Trust land	TQ04	TQ01274920	g2a/c	part-SNCI	SWT/private
SUrk6462	Guildford, Shalford Park	SU94	SU99644862	g3c/fg-m		LA
SUon3819	Guildford, St. Catherines Hill, Artington	SU94	SU99384819	g1a/d		HE
TQrk1024	Guildford, Stoke Meadows & Riverside Park	TQ05	TQ00105124	g3c/fg-m (+f2)	SNCI	LA
SUnt0888	Guildford, The Mount	SU94	SU98084888	g2a/c	SNCI	LA
SUry0992	Guildford, The Mount cemetery	SU94	SU99094892	g2c		CoE
TQns4755	Hackhurst & White Downs, Hackhurst Downs	TQ04	TQ09474855	g2a	SSSI	SCC/SWT, NT
TQws1848	Holmbury St. Mary, Holmdale Meadows	TQ14	TQ11184348	g3c	SNCI	private
TQd 0143	Holmbury St. Mary, St. Mary's churchyard	TQ06	TQ11014443	g1a*	SNCI	CoE
SUow1041	Merrist Wood College, House Meadow	SU95	SU96105341	g3c	SNCI	private
SUs 4028	Merrist Wood College, Low & Water Meadows	SU95	SU96405428	g3c		Activate Learning
TQwn9786	Merrow Downs, Browning Down	TQ04	TQ01974986	g2c	SNCI	LA
SUon1690	Normandy, field E of Henley Park mansion	SU95	SU94165290	g3c		private
SUrm5883	Normandy, field E of Strawberry Farm	SU95	SU93585083	g3c damp		private
SUrd4031	Normandy, fields at Flexford	SU95	SU93405031	g3c	SNCI	private
SU E6330	Normandy, Hunts Hill Farm, wet meadows to E	SU95	SU93635230	g3c damp	part-SNCI, cSNCI	private
SUld1591	Normandy, new cricket field	SU95	SU92155191	g1a/d		PC
TQnd3011	Ockham, fields at Hatchford End	TQ05	TQ08305811	g3c damp		SCC (part)
TQnd5112	Papercourt Marshes NR, dry grassland	TQ04	TQ03515612	g1a	SSSI	private/SWT
TQrm7026	Peaslake, floodplain N of Gasson Farm	TQ04	TQ08704426	g3c		private
TQws7160	Peaslake, Law Brook floodplain meadows	TQ04	TQ08714560	g3c	SNCI	private
SUow9161	Pirbright, Bakersgate Meadow	SU95	SU94915461	g3c damp	SNCI	private
SUrd2691	Pirbright, churchyard	SU95	SU94265591	g1a/d		CoE
SUrm1041	Pirbright, fields around Whites Farm	SU95	SU95105541	g3c damp		private
SUe)7340	Pirbright, fields S of Bowling Hill (composite)	SU95	SU93735440	g1d/g3c	cSNCI	private
SUad4349	Pirbright, fields S of Gole Road	SU95	SU93435649	g1d/g3c	cSNCI	private
SUne8042	Pirbright, fields W of Vapery Lane	SU95	SU93805642	g3c damp		private
SUnd3516	Pirbright, Gapemouth Road, camping ground	SU95	SU92355616	g1a		MoD
SUnd6038	Pirbright, Stoney Castle & Furze Hill camping ground	SU95	SU92605638	g1a/d	SNCI	MoD
SUen6501	Pirbright, village green	SU95	SU94655601	g1a/d	SNCI	PC
SUp)6024	Puttenham Down (Hog's Back scarp)	SU94	SU92604824	g2c	SNCI	SCC/PC
SUds9983	Puttenham, Broomfields	SU94	SU90994483	g1a/d		private
SUne5024	Puttenham, field N of Totford Lane	SU94	SU91504824	g2c		private
SUws1962	Puttenham, Gatwick flood-meadows	SU94	SU91194462	g3c/fg-m		private
SU. 2045	Puttenham, Polo Field, Private	SU94	SU91204745	g1a/d		private
TQt)4141	Ripley, village green (part)	TQ05	TQ05415741	g3c	SNCI	PC
SUow9424	Seale Chalk-pit meadow	SU84	SU89944824	g2a	SNCI	SWT
SUwn7016	Seale, Great Down & Long Down	SU94	SU90704816	g2c		private
SUrk5147	Seale, Hampton Park	SU94	SU90514647	g1a/d, g3c		private
SUds6522	Seale, Seale Common grasslands	SU94	SU90654722	g1a/d		private
TQry3469	Send cemetery	TQ05	TQ02345469	g1a/d		CoE/LA
SUws9314	Shackleford-Peper Harow, Broomfield Meadows	SU94	SU91934414	g3c/fg-m	SNCI	private
SUrd0884	Shackleford, churchyard	SU94	SU94084484	g1a		CoE
SUws6433	Shackleford, Sugarbaker Farm flood-meadows	SU94	SU91644433	g3c/fg-m		private
TQry0517	Shalford cemetery	TQ04	TQ00054717	g3c		LA/CoE
TQon2489	Shalford Common	TQ04	TQ00244689	g1a/d, g3c	SNCI	LA
TQne5437	Shalford, fields by Tillingbourne	TQ04	TQ00544737	g3c/fg-m	part-SNCI	private
TQes6204	Shalford, The Chantries, ex-pastures	TQ04	TQ01624804	g1a/g2c	SNCI	LA
TQs)8890	The Sheepleas, grasslands (inc. Coronation & Summerhouse Meadows)	TQ05	TQ08885190	g2a	SSSI	SCC/SWT
SUNR4557	Unstead Sewage Treatment Works NR	SU94	SU99454557	g3c/fg-m		TW
SUow4047	Wanborough, Inwood Meadow	SU94	SU91404847	g2c	SNCI	private
SUre6081	Wanborough, Manor Farm pasture	SU95	SU93604881	g3c		private
SUe 9551	Wanborough, meadow at Down Place	SU94	SU94954851	g2c		private
TQne9013	West Horsley, Dawes Dene	TQ05	TQ07905113	g2c	SNCI	private
TQne9982	West Horsley, field S of Dawes Dene	TQ05	TQ07995082	g3c damp		private
SUl)7902	Wey Valley Meadows (composite; inc. Broadford flood-meadows-Catteshall)	SU94	SU98794502	g3c/fg-m, g1a (+f2)	part-SSSI	LA, NT, private
TQ')4783	Wisley Common, Park Barn Farm ('field 2')	TQ05	TQ07475983	g1a/d		private

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQTW2963	Wisley, fields E of Wisley STW	TQ05	TQ06295963	g3c		RHS
TQ259844	Wisley, fields S of M25	TQ05	TQ06985944	g1a/d		RHS
SUe)6708	Wood Street, Broadstreet Common (composite)	SU95	SU96675108	g3c	SNCI	SCC
SUen0557	Worplesdon, village green	SU95	SU97055357	g3c		PC
SUrd1645	Wyke, churchyard	SU95	SU92165145	g3c	SNCI	CoE
Guildford/Woking (West Surrey Council) = 4 sites						
TQm)0470	Broad Mead-Newark Lane, floodplain (inc. Papercourt Meadows NR, fields at Papercourt Farm)	TQ05	TQ03045670	g3c/fg-m (+f2)	part-SSSI	SWT/NT, private
TQs)6285	Old Woking-Pyrford, floodplain (inc. Woking Palace meadows)	TQ05	TQ02625685	g3c/fg-m (+f2)	part-SNCI	LA, private
TQws8154	Pyrford-Walsham Lock floodplain meadows	TQ05	TQ04815754	g3c/fg-m (+f2)	part-SNCI	NT
TQws8128	Sutton Place floodplain meadows	TQ15	TQ00815328	g3c/fg-m	part-SNCI, cSNCI	private
Mole Valley (East Surrey Council >2027) = 88 sites						
TQrd5093	Abinger Common, churchyard	TQ14	TQ11504593	g1a/d		CoE
TQow9832	Abinger Common, Goddards Meadow	TQ14	TQ11984532	g1a/d		private
TQrk6270	Abinger Hammer, Abinger Hall park	TQ14	TQ10624770	g3c		private
TQd.6869	Abinger Hammer, meadow S of Kingfisher Farm	TQ04	TQ09744712	g3c		private
TQm 7412	Abinger Hammer, field E of Felday Rd.	TQ04	TQ09684669	g3c damp		private
TQws3186	Ashtead, Rye Meadows	TQ15	TQ17315886	g3c damp		SWT
TQld8910	Ashtead, Woodfield	TQ15	TQ17895910	g3c damp	SNCI	LA
TQub4498	Bookham Common, Isle of Wight grassy scrub	TQ15	TQ12445598	g3c damp	SSSI	NT
TQld6392	Box Hill Farm, large NE field	TQ15	TQ18635092	g2c		NT
TQes7762	Buckland, The Five Acres	TQ25	TQ22775262	g2c		private
TQve5381	Capel, Dairy House Nature Reserve	TQ14	TQ17533981	g3a/c	SNCI	private
TQen1337	Capel, Misbrooks Green	TQ14	TQ18134137	g3c	SNCI	PC
TQrd5275	Capel, St. John-the-Baptist churchyard	TQ14	TQ17524075	g3c*		CoE
TQow1647	Charlwood, Staggers Avon Meadow	TQ24	TQ24164147	g3c		private
TQds0625	Charlwood, Stanhill Court grounds	TQ24	TQ23064225	g3c damp		private
TQow2939	Charlwood, Stanhill Court Meadow	TQ24	TQ23294239	g3a/c	SNCI	private
TQrd9068	Coldharbour, churchyard	TQ14	TQ14904368	g1a/d*		CoE
TQSE1496	Coldharbour, fields to SE	TQ14	TQ15144396	g1a/d		private
TQrm0669	Coldharbour, grassland W of Kitlands Farm	TQ14	TQ15064369	g1a/d		private
TQds2716	Coldharbour, Minnickfold Fields	TQ14	TQ16274316	g3c		private
TQry4804	Dorking Old cemetery	TQ15	TQ17485004	g3c*	SNCI	CoE
TQld4937	Dorking, 10 Acre Field	TQ15	TQ16495037	g2c		private
TQne9536	Dorking, Cotmandene	TQ14	TQ16954936	g3c	SNCI	LA
TQow5237	Forest Green, Latymers Meadow	TQ14	TQ11524037	g3a	cSNCI	private
TQld3164	Forest Green, Lyefield Farm field	TQ14	TQ11314064	g3c		private
TQen2625	Forest Green, village green	TQ14	TQ12264125	g3c	SNCI	PC
TQwn2868	Hackhurst & White Downs, Blatchford Down	TQ14	TQ10284868	g2a	SSSI	private
TQld9505	Hackhurst & White Downs, Chalk Pit Field	TQ14	TQ11954905	g2a	SSSI	private/NT
TQit4066	Hackhurst & White Downs, clay-shoot chalkpit	TQ14	TQ11404866	g2a	SSSI	private
TQs)5300	Hackhurst & White Downs, Denbies Hillside (inc. Secretary's & Lucas Fields)	TQ14-15	TQ13535000	g2a	SSSI	NT
TQys4347	Hackhurst & White Downs, Lamb's Leys	TQ14	TQ12434947	g2a	SSSI	private/NT
TQon7264	Hackhurst & White Downs, Old Plantation	TQ14	TQ10724864	g2a	SSSI	private/NT
TQst7654	Hackhurst & White Downs, Ranges West & East	TQ14	TQ12764954	g2a	SSSI	private/NT
TQld0932	Hackhurst & White Downs, Steers Field	TQ15	TQ14095032	g3c	SSSI	NT
TQst0767	Hackhurst & White Downs, White Down West	TQ14	TQ11074867	g2a	SSSI	private/NT
TQwn1814	Hackhurst & White Downs, Wholcombe Down	TQ14	TQ12184914	g2a	SSSI	private/NT
TQrd5184	Headley, churchyard	TQ25	TQ20515484	g3c		CoE
TQd.4084	Headley, field S of Lodgebottom Rd.	TQ15	TQ19405384	g2c		private
TQow0541	Headley, High Ashurst reservoir meadow	TQ15	TQ19055341	g2c		private
TQld9384	Holmwood Common, former cricket field	TQ14	TQ16934684	g1a/d*	SNCI	NT
TQ247019	Holmwood Common, grassland W of A24	TQ14	TQ16704619	g3c	SNCI	NT
TQ243106	Holmwood Common, grasslands E of A24	TQ14	TQ17314506	g3c*	SNCI	NT
TQer4734	Holmwood Common, Holmwood Corner	TQ14	TQ17474434	g3c*	SNCI	NT
TQld1050	Holmwood Common, old football field	TQ14	TQ17104550	g1a/d*	SNCI	NT
TQll2585	Hookwood, Withy Gill	TQ24	TQ27254285	g3c/fg-m	SNCI	private
TQNR7178	Leatherhead, Common Meadow NR	TQ15	TQ15715678	g3c/fg-m		LA

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQok7269	Leatherhead, field E of A243, N of Rye-brook	TQ15	TQ16725869	g3c		private
TQds6160	Leatherhead, River Lane Fields	TQ15	TQ14615760	g3c/fg-m	SNCI	LA
TQwn5746	Leatherhead, Yarm Way Down	TQ15	TQ17575546	g2c		private
TQns4183	Leigh, Charman's Farm, paddock & lawns	TQ24	TQ21414683	g3c		private
TQld5399	Leigh, cricket field	TQ24	TQ21534699	g3c		PC
TQad7210	Leigh, field E of Smalls Hill Road	TQ24	TQ22724610	g3c		private
TQce3539	Leith Hill, Leith Hill Place	TQ14	TQ13354239	g1a/d*		NT
TQen0448	Little Bookham, village green	TQ15	TQ13045548	g3c		NT
TQrs4980	Mickleham, Bellasis Valley grassland	TQ15	TQ19115278	g2a/c		private
TQnd1178	Mickleham, churchyard	TQ15	TQ17075337	g2c*		CoE
TQrd0737	Mickleham, Fredley Fields	TQ15	TQ17105263	g3c		private
TQds1063	Mickleham, Ashurst Valley & The Junipers	TQ15	TQ18495280	g2c		private
TQon2075	Mickleham, Headley Lane/B2209 junction	TQ15	TQ17205275	g2a*		SCC/private
TQon4694	Mid Holmwood, field W of Holmwood Common	TQ14	TQ16464594	g3a/c	SNCI	private
TQry4243	Mole Gap-Reigate Escarpment, Betchworth Quarry	TQ15	TQ20425143	g2a	SSSI, SAC	SCC
TQe)8784	Mole Gap-Reigate Escarpment, Box Hill grasslands (composite)	TQ15	TQ17875184	g1a/d, g2a, g2c*	SSSI, SAC	NT
TQry1091	Mole Gap-Reigate Escarpment, Brockham Quarry	TQ15	TQ19745109	g2a	SSSI, SAC	SCC/SWT
TQls8146	Mole Gap-Reigate Escarpment, Buckland Hills	TQ25	TQ22815246	g2a/c	SSSI, SAC	private
TQks2154	Mole Gap-Reigate Escarpment, Buckland Limeworks	TQ25	TQ22215253	g2a	SSSI, SAC	private
TQll3820	Mole Gap-Reigate Escarpment, Coneybury Hill	TQ25	TQ23385220	g2a	SSSI, SAC	NT
TQwn2937	Mole Gap-Reigate Escarpment, Fraser Down	TQ25	TQ21295237	g2a/c	SSSI	SWT
TQs)8002	Mole Gap-Reigate Escarpment, Headley Heath (grassland areas)	TQ15	TQ19805302	g2a, g1a*	SSSI, SAC	NT
TQen0300	Mole Gap-Reigate Escarpment, Headley Warren	TQ15	TQ19035400	g2a	SSSI, SAC	private NR
TQll7605	Mole Gap-Reigate Escarpment, Juniper Hill	TQ25	TQ23765205	g2a	SSSI, SAC	NT
TQns2565	Mole Gap-Reigate Escarpment, Mickleham Downs	TQ15	TQ18255365	g2a	SSSI, SAC	NT
TQe)4180	Mole Gap-Reigate Escarpment, Norbury Park grasslands (composite)	TQ15	TQ15415380	g2a/g2c*	SSSI, SAC	SCC/SWT
TQds3874	Newdigate ex-Brickworks NR grasslands	TQ24	TQ20384274	g3a/c	SNCI	SWT
TQse7272	Newdigate, fields W of Tanhouse	TQ14	TQ19724072	g3c damp		private
TQnd2812	Newdigate, Schermuly's grassland	TQ24	TQ20284412	g3c	SNCI	private
TQNR3315	North Holmwood, Inholms Claypit LNR	TQ14	TQ17334715	g3c	SNCI	LA
TQne4610	Ockley, field W of Smugglers Lane	TQ13	TQ15463710	g3a/c	cSNCI	private
TQen6915	Ockley, village green	TQ14	TQ14694015	g3c	cSNCI	PC
TQe)7196	Polesden Lacey restored grasslands; inc. Worzel Bank, Yew Tree Down, Ha-ha Field, Nursery Field (composite)	TQ15	TQ13715196	g2a/c		NT
TQrm6193	Polesden Lacey, fields to N & W of Bagden Farm	TQ15	TQ14615193	g2a/c		NT
TQst2183	Ranmore Common East	TQ15	TQ15215083	g3c		NT
TQC)4641	Ranmore Common, grassy verges (non-SCC)	TQ15	TQ13465041	g3c		NT
TQrd5947	Ranmore, churchyard	TQ15	TQ14595047	g1a/d*		CoE
TQrm5773	Ranmore, fields S of Denbies Farm	TQ15	TQ15575073	g2c		private
TQld6074	Ranmore, Rectory field	TQ15	TQ14605074	g1a/d		private
TQow3263	Wallis Wood NR, meadow	TQ15	TQ12323863	g3a/c	SNCI	SWT
TQSW7387	Westhumble, Chapel Farm, Field to SW	TQ15	TQ15735187	g2c		NT
TQrd7012	Wotton, fields E of St. John's Church & churchyard	TQ14	TQ12704812	g3c*		private
Reigate & Banstead (East Surrey Council >2027) = 26 sites						
TQns6835	Banstead Downs	TQ26	TQ25686135	g2a	SSSI	SCC/BCC
TQ.)1870	Banstead Heath, Chussex Plain East	TQ25	TQ23765381	g3c	SNCI	LA
TQst7681	Banstead Heath, grasslands	TQ25	TQ23675516	g3c	SNCI	BCC
TQds6716	Banstead, Nork Park	TQ25	TQ23675907	g2c	SNCI	LA
TQrk6707	Chipstead Downs (composite, inc. Park & Stagbury Downs, Fames Rough etc.)	TQ25	TQ27185770	g2a/c	SSSI	BCC/LA, SCC
TQds9661	Chipstead, Dene Fields	TQ25	TQ26965761	g2a	SNCI	LA
TQm 5302	Chipstead, fields NW of Outwood Farm	TQ25	TQ26535702	g2c		private
TQrd3639	Chipstead, St. Margaret's churchyard	TQ25	TQ28365639	g1a/d		CoE
TQch7867	Earlswood Common, fields & cricket pitch	TQ24	TQ26784867	g1a/d, g3c	SNCI	LA
TQon2603	Earlswood Common, Whitebushes Common	TQ24	TQ28264803	g3c	SNCI	LA
TQws7193	Hooley Meadows	TQ25	TQ28715693	g2c	SNCI	LA
TQhs3509	Horley, fields W of The Roughs	TQ33	TQ30354209	g3c damp		private
TQrd3506	Kingswood, churchyard	TQ25	TQ22355506	g1a/d		CoE

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQll5412	Mole Gap-Reigate Escarpment, Colley Hill, Queens Park & Reigate Hill	TQ25	TQ24545212	g2a	SSSI, SAC	NT
TQNR6135	Mole Gap-Reigate Escarpment, Dawcombe NR	TQ24-5	TQ21615235	g2a/c*	SSSI, SAC	SWT/private
TQll5829	Mole Gap-Reigate Escarpment, Gatton Park/Wingate Hill	TQ25	TQ26585229	g2a/c	SSSI	NT
TQds7305	Mole Gap-Reigate Escarpment, Reigate Fort & fields	TQ25	TQ25735205	g2a	SSSI, SAC	NT
TQy 7362	Redhill, Redstone cemetery	TQ24	TQ28734962	g1a/d		CoE/LA
TQes0644	Reigate Heath, sports pitches	TQ25	TQ24065044	g1a	SSSI	LA
TQld7186	Reigate, Colley Field	TQ25	TQ24715186	g2c		private
TQow2469	Reigate, Dungate Manor meadow	TQ24	TQ23244969	g1a/d		private
TQll1692	Reigate, pastures at Quarry Hill	TQ25	TQ26165192	g2c		private
TQrd0308	Reigate, St. Mary's churchyard	TQ25	TQ26035008	g1a/d		CoE
TQds9176	South Earlswood, Burnt Oak Farm, Mossy, Anthill & Lime fields	TQ24	TQ26904775	g3c damp		private
TQon7935	South Earlswood, Petridgewood Common	TQ24	TQ27794735	g1a/d, g3c	SNCI	LA
TQrd2856	Walton-on-the-Hill, churchyard	TQ25	TQ24285556	g1a/d		CoE
Runnymede (West Surrey Council >2027) = 17 sites						
TQce0387	Addlestone, Sayes Court public open space	TQ06	TQ05036387	g3c		LA
TQNR9614	Chertsey Meads LNR	TQ06	TQ05966614	g3a/c	SNCI	LA
TQey1826	Chertsey, Barrsbrook Farm, Chertsey	TQ06	TQ03186626	g3c damp	SNCI	LA
TQrm1872	Chertsey, Sandgates car-park	TQ06	TQ03396613	g1a		LA
TQrk3913	Chertsey, Pannells Farm	TQ06	TQ04186572	g3c damp	part-SNCI	private
TQrk2646	Coopers Hill-Runnymede Park	TQ07	TQ00267146	g1a/d	SNCI	private
TQte1881	Englefield Green, Bakeham House estate	TQ06	TQ00186981	g3c damp	cSNCI	private
SUrd7036	Longcross, churchyard	SU96	SU98706536	g1a/d	SNCI	private
TQds9373	Lyne, Hardwick Court Farm fields	TQ06	TQ02936573	g3c damp	SNCI	private
TQrd5670	Lyne, Holy Trinity churchyard	TQ06	TQ01566570	g1a/d		CoE
TQve1975	New Haw, field N of Moated Farm Drive	TQ06	TQ05196375	g3c damp		private
TQrd0949	Ottershaw, churchyard	TQ06	TQ02096349	g1a/d		CoE
TQrk2493	Ottershaw, Homewood Park	TQ06	TQ02246493	g3c		private
TQnd4324	Runnymede & Langham Pond area, inc. Coopers Hill Slopes & Kennedy Memorial grassland	TQ07	TQ00437224	g1a/d, g3c	SNCI/SSSI	NT
TQe)5448	Stroude, fields around Whitehall Farm (composite)	TQ06	TQ00546948	g3a/c damp	cSNCI	private
TQow9804	Thorpe Hay Meadow	TQ07	TQ02987004	g3a	SSSI	SWT
TQow0386	Thorpe, field S of Thorpe Hay Meadow	TQ06	TQ03036986	g3a	cSNCI	private
Spelthorne (West Surrey Council >2027) = 5 sites						
TQow6958	Laleham, Dumsey Meadow	TQ06	TQ05696658	g3a/c	SSSI	LA
TQow9840	Lower Halliford, Thames Meadow	TQ06	TQ08986640	g3c		LA
TQon8277	Staines-upon-Thames, Shortwood Common	TQ07	TQ04827177	g3c	SSSI/SNCI	LA
TQor2610	Staines-upon-Thames, Staines Moor	TQ07	TQ03267310	g1a, g3a/c	SSSI	LA
TQrk6081	Sunbury, Sunbury Park	TQ16	TQ10606881	g3c	SNCI	LA
Surrey Heath (West Surrey Council >2027) = 16 sites						
SUen8778	Bisley, village green	SU95	SU94875978	g1a/d	SNCI	PC
SUld3957	Camberley, Barossa Common, ex-playing field	SU96	SU87396157	g1a/d		MoD
SUrk2465	Camberley, Tekels Park	SU95	SU88245965	g1a/d	SNCI	SHBC
SUne9050	Chobham, Broadford meadows by The Bourne	SU96	SU96906150	g3c/fg-m	SNCI	private
SUrm1222	Chobham, field S of Foxtail Farm	SU96	SU99126322	g3c damp		private
TQrt0809	Chobham, field W of Fairoaks Airport	TQ06	TQ00086209	g3c damp		private
SUws2074	Chobham, Lovelands Farm meadows	SU96	SU96206074	g3c/fg-m	SNCI	private
SUrm6433	Chobham, wet meadows at Larkenshaw Farm	SU96	SU99646233	g3c damp	cSNCI	private
SUry0022	Chobham, wet meadows at Roselands Nursery	SU96	SU96006122	g3c damp	part-SNCI	private
SUrd0812	Frimley, churchyard	SU85	SU88085812	g1a/d		CoE
SUn)3842	Frimley, Hawley Meadows (Surrey section)	SU95	SU86385842	g3c/fg-m	SNCI	LA/BVCP
SUk 0937	Lucas Green, fields N of Trulley Brook	SU96	SU94096037	g1d/g3c	cSNCI	private
SUrd3575	Valley End, St. Saviour's churchyard	SU96	SU95356375	g1a/d	SNCI	CoE
SUow3403	West End, Ralph's Meadow	SU96	SU94346103	g1a/d	SNCI	private
SUrd6937	West End, St. John the Baptist churchyard	SU96	SU94696137	g1a/d	SNCI	CoE
SUn.9764	West End, wet meadows W of Donkey Town/Hook Lane	SU96	SU92976064	g1d/g3c		MoD
Surrey Heath/Woking (West Surrey Council) = 1 site						

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 hab.:	existing site designations:	owner/occupier (if known):
SUws8337	Chobham, Bourne Fields & Young Stroat meadows	SU96	SU99836137	g3c/fg-m	SNCI	private
Tandridge (East Surrey Council >2027) = 73 sites						
TQry3282	Bletchingley cemetery	TQ35	TQ33325082	g1a/d		CoE
TQrd7684	Bletchingley, churchyard	TQ35	TQ32765084	g1a/d		CoE
TQam5166	Bletchingley, field S of Park Ham	TQ35	TQ31515366	g2c		private
TQrs5387	Bletchingley, Quarry Hangers	TQ35	TQ31535387	g2a/c	part-SSSI	SWT/private
TQea8076	Blindley Heath, central grassed area	TQ34	TQ36804476	g3a/c	SSSI	LA/SWT
TQds9528	Burstow, Copper Coin fields	TQ34	TQ31954128	g3c	SNCI	private
TQne9994	Burstow, field to E of Shipleybridge Lane	TQ33	TQ30993994	g3c damp		private
TQod9372	Burstow, fields NE of Bridges Wood	TQ34	TQ30934172	g3c damp	cSNCI	private
TQne8425	Burstow, meadows W of Cogman's Lane	TQ34	TQ32844425	g3c		private
TQry7845	Caterham cemetery	TQ35	TQ33785545	g2a/c		CoE
TQld8207	Caterham, Chalet Field	TQ35	TQ34825407	g2c	SNCI	private
TQnt3316	Caterham, scarp viewpoint	TQ35	TQ34335316	g2a		private
TQrd2631	Caterham, St. John's churchyard	TQ35	TQ34265531	g2a/c		CoE
TQle0010	Chelsham, Nore Hill & Hoving's Hole	TQ35	TQ38005710	g2a/c	part-SNCI	NT/SWT, private
TQrd8809	Chelsham, St. Leonard's churchyard	TQ35	TQ38885909	g2c		CoE
TQws5268	Copthorne, Clay Hall Lane meadows	TQ33	TQ32523968	g3c	SNCI	private
TQad8532	Crowhurst, fields E of Crowhurst Road	TQ34	TQ38854532	g3c/fg-m	cSNCI	private
TQrm7579	Crowhurst, meadow N of Moat Farm	TQ34	TQ37754579	g3c		private
TQrm9857	Dormansland, fields at Hoopers Farm	TQ44	TQ42984257	g3a/c		private
TQll6401	Dormansland, fields E of Old Surrey Hall	TQ43	TQ42644001	g3c damp	cSNCI	private
TQes1273	Dormansland, Lingfield Cernes	TQ44	TQ42124473	g3a/c fg-m	SSSI	private
TQon5399	Farleigh Common	TQ36	TQ36535999	g1a/d	SNCI	PC/SWT
TQnk6073	Farleigh, Chapel Bank	TQ36	TQ38606073	g2a	SNCI	LoWT/BC
TQrd2782	Felbridge, St. Johns churchyard	TQ33	TQ37273982	g1a/d		CoE
TQad5931	Fickleshole, field N of High Hill Road	TQ36	TQ38596031	g2c		private
TQnd0763	Godstone, field(s) to W of Bay Pond	TQ35	TQ35075163	g3c		private
TQd 8177	Godstone, Hilly Field	TQ35	TQ34815177	g1a/d	SNCI	PC
TQrd6832	Horne, churchyard	TQ34	TQ33684432	g3c		CoE
TQod6044	Horne, field S of Bones Wood	TQ34	TQ34604244	g3c damp		private
TQen8617	Hurst Green, Mill Lane Green	TQ35	TQ39865117	g3c		PC
TQnk5441	Kenley Common, Whyteleafe Bank	TQ35	TQ33545841	g2a		CoL
TQon1469	Limpsfield, field W of Itchingwood Common	TQ45	TQ41145069	g3c		private
TQrd6778	Limpsfield, St. Andrews churchyard	TQ45	TQ42675178	g1a/d*		CoE
TQes6917	Lingfield Nature Reserves	TQ34	TQ38694417	g3a/c	SNCI	private
TQrd8978	Lingfield, churchyard	TQ34	TQ38894378	g1a/d		CoE
TQok4889	Lingfield, meadow at Eden Brook	TQ34	TQ39484389	g3c/fg-m		private
TQld5259	Lingfield, Tom's Field	TQ34	TQ38524459	g3c	SNCI	private
TQ251304	Merstham, Quarry Dean Farm, fields N of M25	TQ35	TQ30135404	g2c		private
TQld7004	Newchapel, cricket field	TQ37	TQ35704204	g3c*		PC
TQry9151	Nutfield cemetery	TQ25	TQ29915051	g1a/d		CoE
TQsh1259	Nutfield Marsh	TQ35	TQ30125159	g3c damp	SNCI	PC
TQrd9186	Nutfield, St. Peter & St. Paul churchyard	TQ35	TQ30915086	g1a/d		CoE
TQen7365	Outwood Common, windmill green	TQ34	TQ32734565	g3c	part-SNCI	NT
TQrd9602	Outwood, churchyard	TQ34	TQ31964602	g3c		CoE
TQry0375	Outwood, Harewood Fields & Old Swan Sanctuary	TQ34	TQ33034775	g3c	part-SNCI	NT/private
TQ. 6502	Outwood, Hookhouse Farm, field S of Brickfield Road	TQ34	TQ31654602	g3c		NT/private
TQaw0982	Outwood, Lodge Farm, fields N of Bannister Shaw	TQ34	TQ34094582	g3c damp		NT/private
TQd 3431	Outwood, Stone House Common & Stonehouse Field	TQ34	TQ31344631	g3c	part-SNCI	private
TQen7517	Oxted, Broadham village green	TQ35	TQ38755117	g3c		PC
TQrd0305	Oxted, St. Mary's churchyard	TQ35	TQ39035305	g3c		CoE
TQws4979	South Godstone, Cloverhouse Meadows	TQ34	TQ36494779	g3a/c	SNCI	private
TQd.8531	South Godstone, field E of Tilburstow Hill Rd.	TQ34	TQ35854831	g3c		private
TQod2563	South Nutfield, meadow NE of Philpotes Wood	TQ34	TQ33254963	g3c damp		private
TQrd4515	Tandridge, St. Peter's churchyard	TQ35	TQ37455115	g3c		CoE
TQds1998	Tatsfield, Hill Park Estate fields	TQ45	TQ42195598	g2a/c	cSNCI	SCC/SWT
TQws9476	Tatsfield, Paynesfield Road meadows	TQ45	TQ41945776	g2c		private
TQrd7409	Tatsfield, St. Mary's churchyard	TQ45	TQ41745609	g2a/c		CoE
TQrd9693	Titsey, churchyard	TQ45	TQ40965493	g2a/c		CoE
TQrm8342	Titsey, field SE of Cheverells Farm	TQ35	TQ39835642	g2c		private
TQs)3563	Titsey, Pilgrims Farm field(s)	TQ45	TQ41355563	g2c		private

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQlk1731	Warlingham Court Farm, fields over Chalk	TQ35	TQ34175931	g2c		LA
TQrd6399	Warlingham, churchyard	TQ35	TQ35635899	g2c		CoE
TQce4537	Warlingham, Tippet's Piece	TQ35	TQ35455737	g2a/c		private
TQrd7422	Whyteleafe, All Saints' churchyard	TQ35	TQ33745822	g2c		CoE
TQrk3616	Whyteleafe, Manor Park	TQ35	TQ34365716	g2a/c	SNCI	LA
TQws5620	Woldingham & Oxted Downs, Beau Coin House/(Slines) meadows	TQ35	TQ37565620	g2a/c	SSSI	private
TQll6517	Woldingham & Oxted Downs, Gangers Hill	TQ35	TQ37655417	g2a/c	SSSI	NT
TQke3580	Woldingham & Oxted Downs, South Hawke	TQ35	TQ37355380	g2a	SSSI	NT
TQep7338	Woldingham & Oxted Downs, Whistler's Steep	TQ35	TQ38735438	g2a	SSSI	NT
TQm 0570	Woldingham, Highlands Farm	TQ35	TQ37055770	g2a/c		private
TQll5872	Woldingham, Long Hill	TQ35	TQ36585672	g2a		private
TQds2013	Woldingham, Marden Park grasslands	TQ35	TQ37205413	g2a/c	SSSI	WT
TQrd1590	Woldingham, St. Agatha's churchyard	TQ35	TQ37155490	g2c		CoE
Waverley (West Surrey Council >2027) = 108 sites						
TQ814705	Alfold Crossways, field E of A281	TQ03	TQ04473505	g3c damp		private
TQne5593	Alfold Crossways, field S of Wildwood Lane	TQ03	TQ04553593	g3c damp		private
TQne7577	Alfold, field S of Rosemary Lane	TQ03	TQ02753377	g3c		private
TQch5109	Blackheath, cricket pitch	TQ04	TQ03514609	g1a/d		PC
SUw)3720	Bowlhead Green, fields E of Begley Farm (inc. Soaks Meadow)	SU93	SU90373720	g3c	SNCI	private
TQrm8278	Bramley, field SW of Westland Farm	TQ04	TQ01824378	g3c damp	SNCI	private
TQws3572	Bramley, Snowdenham meadows	TQ04	TQ00354472	g3c		private
TQm 2672	Bramley, wet fields at Bonhurst Farm	TQ04	TQ01264272	g3c damp		private
SUws9016	Charterhouse-Eashing, R. Wey floodplain meadows	SU94	SU94904416	g3c/fg-m(+f2)	SSSI	private
SUrd9838	Chiddingfold, St. Mary's churchyard	SU93	SU95983538	g3c		CoE
SUs 6662	Churt, Borrow House Farm meadows	SU83	SU86663962	g1a/d		private
SUrd3655	Churt, churchyard	SU83	SU85363855	g1a		CoE
SUws7184	Churt, Crosswater Farm meadows	SU83	SU85713984	g1a/d		private
SUen3448	Churt, Redhearn Green	SU83	SU85343848	g1a		PC
TQon0442	Cranleigh Common	TQ03	TQ05043942	g3c	SNCI	LA
TQld7651	Cranleigh, Beryl Harvey Field	TQ03	TQ05763851	g3c	cSNCI	private
TQds6424	Cranleigh, Fernfell Fields	TQ04	TQ06644124	g3c	SNCI	private
TQgh3454	Cranleigh, meadow N of Bowles Rough	TQ03	TQ08343854	g3c damp	cSNCI	private
TQon1434	Cranleigh, Rowly, Smithwood Common	TQ04	TQ05144134	g1a/d	cSNCI	private
SUes2458	Crosswater, Churt, Harold's Hill pastures	SU83	SU85243958	g1a/d		private
SUow9321	Dockenfield, Abbotts Corner meadow	SU84	SU81934021	g3c		private
SUen9710	Dockenfield, Batt's Corner meadow & green	SU84	SU81974110	g3c		private/LA
SUon2792	Dockenfield, Bealeswood Common	SU84	SU82274092	g1a*	SNCI	LA
SUs)8761	Dockenfield, Three Ways meadow(s)	SU84	SU81874061	g3a/c	cSNCI	private/PC
TQch6333	Dunsfold Common, inc. cricket pitch	TQ03	TQ00633633	g3a/c	SNCI	PC/LA
TQne6226	Dunsfold, field E of Rams Lane	TQ03	TQ01623526	g3c		private
TQll5557	Dunsfold, field N of Chapel Hill	TQ03	TQ00553557	g3c damp		private
SUry6517	Eashing cemetery	SU94	SU95654317	g1a/d		CoE
SUll5267	Eashing, flood-meadows at Abbey Mill	SU94	SU94524367	g3c/fg-m (+f2)		private
SUry4753	Elstead (closed) cemetery	SU94	SU90474353	g1a/d		CoE
SUds9277	Elstead, Charles Hill grounds	SU84	SU88924477	g1a/d	SNCI	private
SUR)9194	Elstead, Charleshill (inc. Thundry Meadows NR)	SU84	SU89914394	g3c/fg-m (+f2)	part-SSSI	SWT/private
SUse3955	Elstead, fields W of Westbrook House	SU84	SU89394355	g1a/d		private
SUds9955	Elstead, Hankley Farm fields	SU84	SU88994355	g1a	part-SSSI	private
SUd.2965	Elstead, Meadow at Salena, Thursley Rd.	SU94	SU90294265	g1a/d		private
SUs 2707	Elstead, Polshot Manor meadows	SU84	SU90274407	g3c/fg-m		private
SUow1583	Elstead, Somerset Bridge flood-meadow	SU94	SU92154383	g3c/fg-m (+f2)		private
SUrd3439	Elstead, St. James' churchyard	SU94	SU90344339	g1a/d		CoE
SUrm4388	Elstead, Truxford Wood Farm	SU84	SU89434088	g1a/d		private
SUws9076	Elstead, Westbrook Farm meadows	SU84	SU89904376	g3c/fg-m	part-SNCI	private
SUen0156	Elstead, Westbrook Green	SU94	SU90014356	g1a	SNCI	LA
SUws5326	Elstead, Woodside Farm flood-meadows	SU94	SU91534426	g3c/fg-m		private
SUG 9493	Elstead/Peper Harow, Weyburn Meadows SANG	SU94	SU91944393	g3c/fg-m +f2		Land Trust

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQld4750	Ewhurst Green, village green & cricket field	TQ03	TQ09473950	g3c		LA
TQds5945	Ewhurst, Mullard Space Laboratory fields	TQ14	TQ10594245	g3c damp		UCL
TQrd1348	Ewhurst, St. Peter & St. Paul churchyard	TQ04	TQ09134048	g3c		CoE
SUow4121	Farnham, Bishop's Meadow	SU84	SU83414621	g3c/fg-m	SNCI	PC/private
SUlk8068	Farnham, Compton, field W of Monk's Walk	SU84	SU85804568	g1a/d	SNCI	private
SUds1195	Farnham, Farnham Park grasslands	SU84	SU84114795	g1a/d	SNCI	LA
SUey9426	Farnham, fields E & W of Waverley Abbey	SU84	SU86944526	g3c/fg-m	cSNCI	private/HE
SUch2219	Farnham, Snayleslynch	SU84	SU85224719	g3c/fg-m	SNCI	LA
SUds5524	Frensham, Frensham Manor Fields	SU84	SU83554024	g3c/fg-m	SNCI	private
SUrd1547	Frensham, St. Mary-the-Virgin churchyard	SU84	SU84154147	g3c		CoE
SUry2065	Godalming, Nightingale Rd. cemetery	SU94	SU97204465	g1a/d		PC/CoE
SUws7432	Godalming, Overgone & Lammas Meadows	SU94	SU97744432	g3c/fg-m	SNCI	LA
TQrd3065	Grafham, St. Andrew's churchyard	TQ04	TQ02304165	g3c		CoE
TQws5438	Grafham-Cranleigh, East Whipley Farm flood-meadows	TQ04	TQ03544138	g3c/fg-m		private
SUrd7280	Grayswood, All Saints churchyard	SU93	SU91723480	g1a/d		CoE
SUld1809	Grayswood, Clammer Hill Field	SU93	SU92183409	g1a/d		private
SUen7871	Grayswood, cricket field & village green	SU93	SU91783471	g1a/d		LA/NT
SUte7998	Grayswood, fields at Grayswood Hill Estate	SU93	SU91793398	g3c		private
TQrd1753	Hascombe, churchyard	TQ03	TQ00173953	g1a/d		CoE
SUne7820	Hascombe, field N of Markwick Lane	SU93	SU99783820	g1a/d		private
SUon6755	Haslemere, Almshouse Common	SU93	SU91673255	g3a/c		NT
SUrm5988	Haslemere, fields E of Stedlands Farm	SU83	SU90593188	g3c		NT
SUrk4289	Haslemere, Lythe Hill Park	SU93	SU91423189	g1a/d		private
SUws3802	Haslemere, Meadfields Farm meadows	SU93	SU91383302	g3c		private
SUws2501	Haslemere, Nutcombe Lane meadows	SU83	SU88253401	g3c damp		private
SUry6273	Haslemere, Sunvale Cemetery	SU83	SU87533272	g1a/d		LA
SUws0756	Haslemere, Valewood meadows	SU83	SU91073156	g3c/damp		NT
SUs 4233	Milford, Kennel Moor meadows	SU94	SU93424233	g3c damp	cSNCI	private
SUws6812	Milford, Mistlebrooks & Borough Farm Meadows	SU94	SU92684212	g3c damp	SNCI	private
SUws0061	Millbridge, Burtleys meadows	SU84	SU85004261	g1a/d		private
SUld5515	Millbridge, Frensham cricket field	SU84	SU84554215	g1a/d		PC
SUow9720	Millbridge, Pierrepont Farm flood-meadow	SU84	SU85974220	g3c/fg-m		CRT
SU. 5600	Millbridge, S of Millbridge Rec.	SU84	SU84564200	g1a/d		PC
SUon1123	Millbridge, Shortfield Common	SU84	SU84114223	g1a/d	SNCI	LA
SU5 6099	Millbridge, Thursley,Hankley & Frensham Commons, Unit 5	SU84	SU85604199	g3c/fg-m (+f2)	SSSI	CRT
SUrk0626	Peper Harow Park	SU94	SU93064426	g1a/d		private
SUws9797	Peper Harow Park flood-meadows	SU94	SU92974397	g3c/fg-m		private
SUsh6254	Peper Harow Park, marsh	SU94	SU93624354	g3c/fg-m	SNCI	private
SUws5938	Rowledge, Frensham Vale Farm meadows	SU84	SU83594338	g3c		private
SUow6141	Rowledge, Oast House Meadow	SU84	SU82614241	g3a/c	SNCI	private
TQon2283	Rowly, Run Common	TQ04	TQ03224183	g3c		private
TQry3746	Shamley Green, churchyard & cemetery	TQ04	TQ03374346	g1a/d		CoE
TQen1784	Shamley Green, village green	TQ04	TQ03174384	g3c		PC
SUws1954	Spreakley, flood-meadows	SU84	SU84194154	g3c/fg-m		private
SUow1076	Spreakley, Little Oakley meadow	SU84	SU82104176	g3c		private
SUld8272	The Bourne, cricket field	SU84	SU84824472	g1a/d		LA
SUss5253	The Sands, field at Littleworth Cross	SU84	SU89524553	g1a/d		private
SUnd9387	Thursley, cricket field & recreation ground	SU83	SU89933987	g1a		PC
SUrd1037	Thursley, St. Michael & All Angels churchyard	SU93	SU90103937	g1a/d		CoE
SUld5553	Thursley-Frensham Commons, Parish Field	SU84	SU90554053	g1a/d	SSSI	NE/PC
SUrd2915	Tilford, churchyard	SU84	SU87294315	g1a		CoE
SUon5989	Tilford, Field E of Tankersford Common	SU84	SU86594289	g1a	cSNCI	RSPB
SUne3892	Tilford, field S of Squires Hill Lane	SU84	SU87384392	g1a/d		private
SUe 0967	Tilford, Stonehills campsite	SU84	SU87094267	g1a/d	cSNCI	SCC/private
SUws3854	Tilford, Tankersford R. Wey watermeadows	SU84	SU86384254	g3c/fg-m		private
SUws4715	Tilford, Whitmead flood-meadows	SU84	SU88474315	g3c/fg-m (+f2)	SNCI	private
SUws6015	Winkworth Arboretum, meadows	SU94	SU99604115	g3a/c	SNCI	NT
TQry7763	Wonersh & Blackheath cemetery	TQ04	TQ02774563	g3c		CoE
TQea2976	Wonersh, Chinthurst Hill summit area	TQ04	TQ01294576	g1a		SCC
TQll4432	Wonersh, field SE of Chinthurst Hill	TQ04	TQ01444532	g3c		private
TQow9509	Wonersh, Lostiford meadow	TQ04	TQ02954509	g3c damp	SNCI	PC

GI_ID:	site name:	hectad (10^2km grid square):	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
TQen8053	Wonersh, village green	TQ04	TQ01804553	g3c		PC
SUs)2186	Woolmer Hill, Kingfisher Farm field(s)	SU83	SU87213286	g3c damp	SNCI	private
SUry3426	Wrecclesham cemetery	SU84	SU83344526	g1a/d		CoE
SUs 3231	Wrecclesham, Willey Mill flood-meadows	SU84	SU82324531	g3c/fg-m		private
Woking (West Surrey Council >2027) = 9 sites						
SUd)6953	Brookwood Cemetery grasslands (non-heathland)	SU95	SU95695653	g1a/d*	SNCI	CoE/LA
SUNG4952	Brookwood, Brookwood Meadows SANG	SU95	SU95495752	g3c damp	SNCI	LA
TQws5490	Byfleet, Common Meadows	TQ05	TQ06545990	g3c/fg-m	SNCI	LA
TQt)9103	Byfleet, Manor Farm NR (part)	TQ06	TQ06916003	g3c/fg-m	SNCI	SWT
TQrd3230	Byfleet, St. Mary's churchyard	TQ06	TQ06326030	g1a/d		CoE
SUne7395	Knaphill, field S of Carthouse Lane	SU95	SU97735995	g3c damp		private
SUds4164	Mimbridge, Millbrook Animal Centre fields	SU96	SU97416064	g3c damp (+f2)	SNCI	RSPCA
TQge5722	West Byfleet, Wey floodplain S of Murray's Bridge	TQ06	TQ05576027	g3c/fg-m		NT
SUs 0029	Worplesdon-Mayford, Hoe Stream meadows	SU95	SU98005529	g3c/fg-m	SNCI	private
Total = **460** site units					* = **grassland fungi site** (≥**10 'CHEGD' spp. recorded**)	

Abbreviata: BC=Butterfly Conservation Surrey & SW London Branch; BCC=Banstead Commons Conservators; BVCP=Blackwater Valley Countryside Project; CCT=Churches Conservation Trust; CoE=Church of England; CoL=City of London; CRT=Countryside Regeneration Trust; f2=UK HAB Freshwater group '2'; fg-m=floodplain grazing-marsh; HE=Historic England; LA=Local Authority (Borough/District); LNR=Local Nature Reserve; LoWT=London Wildlife Trust; MoD=Ministry of Defence; NH=National Highways; NHS=National Health Service; NR=nature reserve; NRA=National Rifle Association; NT=National Trust; PC=local Parish/Town Council; RHS=Royal Horticultural Soc.; RSPCA=Royal Soc. for Prevention of Cruelty to Animals; SAC=Special Area of Conservation; SCC=Surrey County Council; SNCI=Site of Nature Conservation Importance; SSEW=Sutton & South-East Water; SSSI=Site of Special Scientific Interest; STW=sewage treatment works; SWT=Surrey Wildlife Trust; District; TW=Thames Water; UCL=Univ. College London; WT=Woodland Trust.

7. References & further reading

Blakesley, D. & Buckley P. (2016). Grassland Restoration and Management (Conservation Handbooks). Pelagic Publishing

Lowland Grassland Management Handbook (2nd edition). Natural England 1999

Kirby, P. (2013). Habitat management for invertebrates: a practical handbook. Pelagic Publishing

Sankey, P A et al. eds. (2024). Surrey Rare Plant Register: the rare, scarce & declining plants of Surrey. Surrey Botanical Society

Surrey Nature Recovery – Consultation Draft, July 2025. Surrey County Council

Waite, M. (2017). The State of Surrey's Nature. Surrey Nature Partnership

Walker, K.J., Stroh, P.A., Roy, D.B., Burkmar, R.J. & Prescott, O.L. (2003). *Britain's Changing Flora. A Summary of the Results of Plant Atlas 2020.* BSBI

Splendid Waxcap *Hygrocybe splendidissima*

Annex 1. Surrey golf courses supporting significant grassland biodiversity interest

borough	Site	Grid Ref.	UK Hab 2.0
Reigate & Banstead	Banstead Downs (SSSI)	TQ250609	g2c/g1a
Mole Valley	Beaverbrook (Cherkley)	TQ181548	g2c
Mole Valley	Betchworth Park	TQ187497	g1d/3c
Tandridge	Bletchingley (SNCI)	TQ331513	g3c
Surrey Heath	Camberley Heath (SNCI)	SU892596	g1a/c
Reigate & Banstead	Chipstead (SNCI)	TQ280584	g2c
Guildford	Clandon (inc. SNCI)	TQ036505	g2c
Elmbridge	Ditton Common (SNCI)	TQ148660	g1a/g3c
Mole Valley	Dorking	TQ176487	g1d
Guildford	Effingham (SNCI)	TQ114525	g2a/c
Epsom & Ewell	Epsom Downs (SNCI)	TQ222588	g2a/c
Tandridge	Farleigh Court	TQ375612	g2c
Waverley	Farnham (SNCI)	SU884466	g2c
Waverley	Farnham Park (SNCI)	SU838477	g1a/d
Runnymede	Fox Hills	TQ010653	g1a/d
Mole Valley	Gatton Manor	TQ128391	g3c
Guildford	Guildford (SNCI)	TQ030498	g2a/c, g1d
Waverley	Hankley (SSSI)	SU880421	g1a/d
Waverley	Hindhead (SNCI)	SU863373	g1a/d
Woking	Hook Heath (SNCI)	SU977570	g1a/d
Runnymede	Laleham Burway (SNCI) [closed]	TQ045681	g3c/g1d
Guildford	Merrist Wood	SU952531	g3c
Woking	New Zealand (SNCI)	TQ023611	g1a/d
Surrey Heath	Pine Ridge	SU904586	g1a/d
Waverley	Puttenham	SU943476	g1a/d
Woking	Pyrford	TQ052587	g1d/g3c
Runnymede	Queenwood (SNCI)	TQ008639	g1a/d
Reigate & Banstead	Reigate Heath (SSSI)	TQ236501	g1a
Reigate & Banstead	Reigate Hill	TQ283533	g2c
Elmbridge	St George's Hill (SNCI)	TQ078619	g1d/g3c
Surrey Heath	Sunningdale (SNCI)	SU951656	g1d/g3c
Surrey Heath	Sunningdale Ladies (SNCI)	SU948660	g1d/g3c
Tandridge	Surrey National	TQ319558	g2c
Tandridge	Tandridge	TQ377515	g3c/g1d
Woking	Twisted Stone [closed]	TQ052587	g1d/g3c
Mole Valley	Tyrrells Wood	TQ187547	g2c
Reigate & Banstead	Walton Heath (SNCI)	TQ230536	g1a/g2c
Runnymede	Wentworth (part-Chobham Common SSSI/SNCI)	SU970664	g1a/d
Woking	West Byfleet	TQ033603	g1d
Woking	West Hill (SNCI)	SU965567	g1d/g3c
Waverley	Wildwood (Alfold) Golf Couse [closed]	TQ048355	g3c
Tandridge	Woldingham [closed]	TQ360576	g2c
Epsom & Ewell	Woodcote Park	TQ206584	g2c
Guildford	Worplesdon	SU966556	g1d/g3c

Annex 2a. Grassland axiophytes in Surrey

Taxon	Vernacular	UK Hab Classification 2.0	UK Red-List[1]
Achillea ptarmica	Sneezewort	g3a/g3c	
Agrimonia procera	Fragrant Agrimony	g3a/g3c	
Agrostis canina	Velvet Bent	g1d (damp)	
Agrostis curtisii	Bristle Bent	g1a/g1d (dry)	
Agrostis vinealis	Brown Bent	g1d (dry)	
Aira caryophyllea	Silver Hair-grass	g1a/g1d (dry)	**VU**
Aira praecox	Early Hair-grass	g1a/g1d (dry)	**VU**
Allium oleraceum	Field Garlic	g1a/g1d (dry)	
Anacamptis pyramidalis	Pyramidal Orchid	g2a/g2c	
Angelica sylvestris	Wild Angelica	wide range (damp)	
Anthoxanthum odoratum	Sweet Vernal-grass	wide range	
Anthriscus caucalis	Bur Chervil	g1a/g1d (dry)	
Anthyllis vulneraria	Kidney Vetch	g2a/g2c	**NT**
Aphanes australis	Slender Parsley-piert	g1a/g1d (dry)	
Arabis hirsuta	Hairy Rock-cress	g2a	**VU**
Arenaria serpyllifolia	Thyme-leaved Sandwort	g2a/g2c	
Asperula cynanchica	Squinancywort	g2a/g2c	**VU**
Avenella flexuosa	Wavy Hair-grass	g1a/g1d (dry)	
Avenula pubescens	Downy Oat-grass	g2a/g2c/g3a (Thames Valley)	
Betonica officinalis	Betony	g3a/g3c	**VU**
Bistorta officinalis	Common Bistort	g3a/g3c	
Blackstonia perfoliata	Yellow-wort	g2a/g2c	
Briza media	Quaking-grass	g2a/g2c	**VU**
Bromopsis erecta	Upright Brome	g2a/g2c	
Bromus commutatus	Meadow Brome	g3a/g3c	
Bromus racemosus	Smooth Brome	wide range	
Campanula glomerata	Clustered Bellflower	g2a	
Campanula rotundifolia	Harebell	g1a/g1d, g2a	
Cardamine pratensis	Lady's Smock	g3c (damp)	
Carduus nutans	Musk Thistle	g2a/g2c	
Carex acuta	Slender Tufted-sedge	g3c, g1d (damp)	
Carex arenaria	Sand Sedge	g1a	
Carex caryophyllea	Spring-sedge	g1a, g2a	**NT**
Carex demissa	Common Yellow-sedge	g1d/g3c (damp)	
Carex disticha	Brown Sedge	g3c (damp)	
Carex filiformis	Downy-fruited Sedge	g3a/g3c (Thames Valley)	
Carex flacca	Glaucous Sedge	wide range	
Carex leporina	Oval Sedge	wide range	
Carex muricata subsp. pairae	Prickly Sedge	g1a/g1d (dry)	
Carex nigra	Common Sedge	wide range (damp)	
Carex panicea	Carnation Sedge	g1d/g3c (damp)	**NT**
Carex pilulifera	Pill Sedge	g1a/g1d (dry)	
Carex spicata	Spiked Sedge	g2c/g3c	
Carlina vulgaris	Carline Thistle	g2a/g2c	
Centaurea debeauxii	Chalk Knapweed	g2a/g2c	
Centaurea scabiosa	Greater Knapweed	g2a/g2c	
Centaurium erythraea	Common Centaury	wide range (dry)	
Centaurium pulchellum	Lesser Centaury	g1a/g2a	
Cerastium arvense	Field Mouse-ear	g1a	**EN**
Cerastium semidecandrum	Little Mouse-ear	g1a/g1d (dry)	**NT**
Chamaemelum nobile	Chamomile	g1a/g3c	**EN**
Cirsium acaule	Dwarf Thistle	g2a/g2c	**VU**
Cirsium dissectum	Meadow Thistle	g1d (damp)	**EN**
Cirsium palustre	Marsh thistle	g1d/g3c (damp)	
Clinopodium acinos	Basil Thyme	g2a/g2c	**VU**

Clinopodium vulgare	Wild Basil	g2a/g2c	
Conopodium majus	Pignut	wide range (pH <6)	
Crepis biennis	Rough Hawk's-beard	g3a/g3c/g2c	
Cruciata laevipes	Crosswort	g2a/g2c/g3c	
Cuscuta epithymum	Common Dodder	g1a, g2a	
Cynoglossum officinale	Hound's-tongue	wide range (dry)	
Cynosurus cristatus	Crested Dog's-tail	wide range	
Dactylorhiza fuchsii	Common Spotted-orchid	wide range	
Dactylorhiza maculata	Heath Spotted-orchid	g1a/g1d	
Dactylorhiza praetermissa	Southern Marsh-orchid	g3c (damp)	
Danthonia decumbens	Heath-grass	g1a, g2a	**NT**
Daucus carota	Carrot	wide range	
Dianthus deltoides	Maiden Pink	g1a/g1d (dry)	**NT**
Epilobium palustre	Marsh Willowherb	g3c (damp)	
Equisetum palustre	Marsh Horsetail	g3c (damp)	
Erigeron acris	Blue Fleabane	wide range (pH >6)	
Erodium cicutarium	Common Stork's-bill	g1a/g1d (dry)	
Erophila glabrescens	Glaborous Whitlow-grass	g1a	
Erophila verna	Common Whitlow-grass	wide range (pH <6)	
Euphrasia nemorosa	Common Eyebright	wide range	
Euphrasia officinalis ssp. anglica	English Eyebright	g1a/g1d (dry)	
Euphrasia pseudokerneri	Chalk Eyebright	g2a	**NT**
Festuca filiformis	Fine-leaved Sheep's Fescue	g1a	
Festuca ovina	Sheep's-fescue	wide range	
Filago vulgaris	Common Cudweed	g1a/g1d (dry)	
Filipendula ulmaria	Meadowsweet	g3c (damp)	
Filipendula vulgaris	Dropwort	g2a, g3c (pH >6)	
Galium palustre	Marsh Bedstraw	wide range (damp)	
Galium pumilum	Slender Bedstraw	g2a	
Galium saxatile	Heath Bedstraw	g1a/g1d (dry)	
Galium uliginosum	Fen Bedstraw	g1d (damp)	
Galium verum	Lady's Bedstraw	wide range	
Genista anglica	Petty Whin	g1a/g1d/g3c	**EN**
Genista tinctoria	Dyer's Greenweed	g3a/g3c	**EN**
Gentianella amarella	Autumn Gentian	g2a/g2c	
Geranium columbinum	Long-stalked Crane's-bill	g2a	**VU**
Geranium pratense	Meadow Crane's-bill	g3a/g3c (Thames Valley)	
Gymnadenia conopsea	Chalk Fragrant Orchid	g2a	**NT**
Helianthemum nummularium	Common Rock-rose	g2a/g2c	**VU**
Helichtochloa pratensis	Meadow Oat-grass	wide range	
Herminium monorchis	Musk Orchid	g2a	**EN**
Himantoglossum hircinum	Lizard Orchid	g2a/g2c	
Hippocrepis comosa	Horseshoe Vetch	g2a/g2c	**VU**
Hordeum secalinum	Meadow Barley	g3a/g3c	
Hydrocotyle vulgaris	Marsh Pennywort	g1a (damp)	**VU**
Hypericum hirsutum	Hairy St. John's-wort	g2a/g2c	
Hypericum humifusum	Trailing St. John's-wort	g1a/g1d/g3c (dry)	
Hypericum maculatum	Imperforate St John's-wort	g3a/g3c	
Hypericum montanum	Pale St. John's-wort	g2a	**VU**
Hypericum tetrapterum	Square-stalked St. John's-wort	g3c/g1d (damp)	
Hypochaeris glabra	Smooth Cat's-ear	g1a	
Hypochaeris radicata	Common Cat's-ear	wide range (pH <6)	
Inula conyzae	Ploughman's-spikenard	g2a/g2c	
Isolepis setacea	Bristle Club-rush	g1d (damp)	
Jacobaea aquaticus	Marsh Ragwort	g3c (damp)	
Jacobaea erucifolius	Hoary Ragwort	wide range	
Jasione montana	Sheep's-bit	g1a/g1d (dry)	**VU**
Juncus acutiflorus	Sharp-flowered Rush	g1d/g3c (damp)	
Juncus articulatus	Jointed Rush	g1d/g3c (damp)	

Juncus conglomeratus	Compact Rush	g3c/g1d (damp)	
Juncus squarrosus	Heath Rush	g1d (damp)	
Knautia arvensis	Field Scabious	wide range (pH >6)	
Koeleria macrantha	Crested Hair-grass	g2a	**NT**
Lathyrus aphaca	Yellow Vetchling	g3c*	**VU**
Lathyrus linifolius	Bitter-vetch	g1d, g3a/c	**VU**
Lathyrus nissolia	Grass Vetchling	wide range	
Lathyrus pratensis	Meadow Vetchling	wide range	
Leontodon hispidus	Rough Hawkbit	g2a/g2c	
Leontodon saxatilis	Lesser Hawkbit	wide range	
Linum catharticum	Fairy Flax	g2a/g2c	
Linum linoides	Allseed	g1a/g1d (damp)	
Lithospermum officinale	Common Gromwell	g2c	
Logfia minima	Small Cudweed	g1a/g1d (dry)	
Lotus pedunculatus	Greater Bird's-foot-trefoil	g1d/g3c (damp)	
Luzula campestris	Field Wood-rush	wide range	
Luzula multiflora	Heath Wood-rush	g1a/g1d (dry)	
Lysimachia tenella	Bog Pimpernel	g1d (damp)	
Mentha arvensis	Corn Mint	wide range	**VU**
Moenchia erecta	Upright Chickweed	g1a	**VU**
Montia fontana	Blinks	g1a	
Myosotis discolor	Changing Forget-me-not	g1a/g1d (dry)	
Myosotis ramosissima	Early Forget-me-not	g1a	
Nardus stricta	Mat-grass	g1a	
Neottia ovata	Common Twayblade	g2a/g2c	
Oenanthe fistulosa	Tubular Water-dropwort	g3c (damp)	**EN**
Oenanthe pimpinelloides	Corky-fruited Water-dropwort	g3a/g3c	
Oenanthe silaifolia	Narrow-leaved Water-dropwort	g3c (damp)	
Omalotheca sylvaticum	Heath Cudweed	g1a/g1d (dry)	**VU**
Ononis repens	Common Restharrow	g2a/g2c	**VU**
Ophioglossum vulgatum	Adder's-tongue Fern	wide range (pH <6)	
Ophrys apifera	Bee Orchid	wide range (pH >6)	
Orchis anthropophora	Man Orchid	g2a	**VU**
Origanum vulgare	Wild Marjoram	g2a/g2c	
Ornithopus perpusillus	Bird's-foot	g1a	
Pedicularis sylvatica	Lousewort	g1a/g1d	**NT**
Phyteuma orbiculare	Round-headed Rampion	g2a	
Picris hieracioides	Hawkweed Oxtongue	wide range (pH >6)	
Pilosella officinarum	Mouse-ear-hawkweed	wide range	
Pimpinella saxifraga	Burnet-saxifrage	wide range	**NT**
Plantago coronopus	Buck's-horn Plantain	g1a/g1d (dry)	
Plantago media	Hoary Plantain	g1a, g2a	
Poa angustifolia	Narrow-leaved Meadow-grass	g2a/g2c/g1a	
Poa bulbosa	Bulbous Meadow-grass	g1a	
Poa humilis	Spreading Meadow-grass	wide range (pH <6)	
Poa pratensis	Smooth Meadow-grass	wide range	
Polygala calcarea	Chalk Milkwort	g2a	**VU**
Polygala vulgaris	Common Milkwort	g2a/g2c	**VU**
Potentilla anglica	Trailing Tormentil	g1a/g1d (dry)	
Potentilla argentea	Hoary Cinquefoil	g1a	
Potentilla erecta	Tormentil	g1a/g1d/g3c (dry)	
Poterium sanguisorba	Salad Burnet	g2a/g2c	**VU**
Primula veris	Cowslip	g2a/g2c	
Ranunculus bulbosus	Bulbous Buttercup	wide range	
Rhinanthus minor	Yellow-rattle	wide range	
Rumex acetosa	Common Sorrel	wide range	
Rumex acetosella	Sheep's Sorrel	g1a/g1d (dry)	**NT**
Rumex pulcher	Fiddle Dock	g1a/g1d (dry)	**NT**

Sagina nodosa	Knotted Pearlwort	g1a	**VU**
Saxifraga granulata	Meadow Saxifrage	g1a/g1d (dry)	**VU**
Scabiosa columbaria	Small Scabious	g2a/g2c	
Schedonorus pratensis	Meadow Fescue	wide range	**VU**
Scleranthus annuus	Annual Knawel	g1a	**EN**/DD
Serratula tinctoria	Saw-wort	g1a/g1d/g3c	**VU**
Silaum silaus	Pepper-saxifrage	g3a/g3c	**EN**
Silene flos-cuculi	Ragged Robin	g3c (damp)	
Solidago virgaurea	Goldenrod	g1a	**VU**
Spergularia rubra	Sand Spurrey	g1a	
Spiranthes spiralis	Autumn Lady's-tresses	g2a	
Stachys palustris	Marsh Woundwort	g3c (damp)	
Stellaria alsine	Bog Stitchwort	g1d/g3c (damp)	
Stellaria graminea	Lesser Stitchwort	wide range	
Stellaria pallida	Lesser Chickweed	g1a/g1d (dry)	
Stellaria palustris	Marsh Stitchwort	g3c (damp)	**EN**
Succisa pratensis	Devil's-bit Scabious	wide range (pH <6)	**VU**
Teesdalia nudicaulis	Shepherd's Cress	g1a	**NT**
Thesium humifusum	Bastard Toadflax	g2a	**VU**
Thalictrum flavum	Common Meadow Rue	g3a/g3c	
Thymus drucei	Wild Thyme	g1a, g2a	**NT**
Thymus pulegioides	Large Thyme	g2a	**VU**
Trifolium arvense	Hare's-foot Clover	g1a/g1d (dry)	**VU**
Trifolium fragiferum	Strawberry Clover	g3c	**EN**
Trifolium medium	Zigzag Clover	g3a/g3c/g1d	**VU**
Trifolium micranthum	Slender Trefoil	g1a	
Trifolium ornithopodioides	Bird's-foot Clover	g1a	
Trifolium striatum	Knotted Clover	g1a/g1d (dry)	
Trifolium subterraneum	Subterranean Clover	g1a	
Trisetum flavescens	Yellow Oat-grass	wide range (pH >6)	
Turritis glabra	Tower Mustard	g1a	**NT**
Veronica officinalis	Heath Speedwell	g1a/g3c (dry)	
Vicia lathyroides	Spring Vetch	g1a	**VU**
Vicia sativa subsp. nigra	Narrow-leaved Vetch	g1a/g1d (dry)	
Viola canina	Heath Dog-violet	g1a	**VU**
Viola hirta	Hairy Violet	g2a/c	

pH >6=calcareous inclination; <6=acidic inclination *naturalised in admin. Surrey

[1] *A new Vascular Plant Red list for Great Britain 2025* = 'GBRL 2025'; IUCN Conservation categories: Endangered = EN; Vulnerable = VU; Near Threatened = NT

Annex 2b. CHEGD fungi species in UK

Clavarioids (corals, spindles & clubs - ***Clavaria***, ***Clavulinopsis***, ***Ramariopsis***)

Clavaria amoenoides
Clavaria atroumbrina
Clavaria fragilis
Clavaria fumosa
Clavaria greletii
Clavaria incarnata
Clavaria straminea
Clavaria tenuipes
Clavaria zollingeri HDI
Clavulinopsis candida
Clavulinopsis corniculata
Clavulinopsis fusiformis
Clavulinopsis helvola
Clavulinopsis laeticolor
Clavulinopsis luteoalba
Clavulinopsis umbrinella
Ramariopsis crocea
Ramariopsis kunzei
Ramariopsis luteonana
Ramariopsis minutula
Ramariopsis pulchella
Ramariopsis subtilis
Ramariopsis tenuiramosa

(Waxcaps, including ***Hygrocybe*** sens.lat. ***Camarophyllus***, ***Cuphophyllus***, ***Gliophorus***, ***Porpolomopsis***)

Chromosera citrinopallida
Cuphophyllus aurantius
Cuphophyllus canescens
Cuphophyllus colemannianus HDI
Cuphophyllus flavipes
Cuphophyllus fornicates
Cuphophyllus lacmus HDI
Cuphophyllus lepidotus
Cuphophyllus pratensis
Cuphophyllus radiatus
Cuphophyllus russocoriaceus
Cuphophyllus virgineus
Gliophorus europerplexus
Gliophorus irrigatus
Gliophorus laetus
Gliophorus perplexus aff.
Gliophorus psittacinus
Gliophorus reginae
Gliophorus vitellinus
Hygrocybe acutoconica
Hygrocybe acutoconica* var. *konradii
Hygrocybe angustifolia
Hygrocybe aurantiosplendens HDI
Hygrocybe calciphila
Hygrocybe cantharellus
Hygrocybe ceracea
Hygrocybe chlorophana
Hygrocybe citrinovirens HDI
Hygrocybe coccinea
Hygrocybe conica
Hygrocybe conicoides
Hygrocybe constrictospora
Hygrocybe glutinipes
Hygrocybe helobia
Hygrocybe insipida
Hygrocybe intermedia HDI

Hygrocybe marchii
Hygrocybe miniata
Hygrocybe monteverdae
Hygrocybe mucronella
Hygrocybe olivaceonigra
Hygrocybe phaeococcinea
Hygrocybe punicea
Hygrocybe quieta
Hygrocybe reidii
Hygrocybe spadicea HDI
Hygrocybe splendidissima HDI
Hygrocybe subpapillata HDI
Hygrocybe subradiatus
Hygrocybe substrangulata
Hygrocybe turunda HDI
Hygrocybe xanthochroa
Neohygrocybe ingrata HDI
Neohygrocybe nitrata HDI
Neohygrocybe ovina HDI
Porpolomopsis calyptriformis HDI

(Pinkgills & allies - ***Entoloma*** spp.)
Entoloma ameides
Entoloma anatinum
Entoloma argentiostriatum
Entoloma asprellum
Entoloma atrocoeruleum
Entoloma atromadidum
Entoloma atromarginatum
Entoloma bloxamii agg. HDI
Entoloma caeruleopolitum
Entoloma caeruleum
Entoloma caesiocinctum
Entoloma carneogriseum
Entoloma catalaunicum
Entoloma chalybeum/lazulinum
Entoloma clandestinum
Entoloma cocles
Entoloma corvinum
Entoloma cyaneoviridescens
Entoloma excentricum
Entoloma exile
Entoloma favrei
Entoloma fernandae
Entoloma formosum
Entoloma fuscotomentosum
Entoloma griseocyaneum
Entoloma griseorubidum
Entoloma henrici
Entoloma incanum
Entoloma indutoides
Entoloma infula
Entoloma insidiosum
Entoloma kervernii
Entoloma kuehnerianum
Entoloma lampropus
Entoloma langei
Entoloma lividocyanulum
Entoloma longistriatum
Entoloma madidum
Entoloma mougeotii
Entoloma nigroviolaceum
Entoloma occultopigmentatum
Entoloma ochromicaceum

Entoloma ortonii
Entoloma ochreoprunuloides
Entoloma papillatum
Entoloma poliopus
Entoloma porphyrophaeum
Entoloma prunuloides
Entoloma pseudoturci
Entoloma roseum
Entoloma saccharioleus
Entoloma sodale
Entoloma solstitiale
Entoloma turci
Entoloma xanthocaulon
Entoloma xanthochroum

(Earthtongues & allies, including ***Geoglossum***, ***Glutinoglossum***, ***Microglossum***, ***Trichoglossum***)
Geoglossum atropurpureum s. str.
Geoglossum barlae
Geoglossum cookeianum
Geoglossum fallax
Geoglossum nigritum
Geoglossum simile
Geoglossum starbaeckii
Geoglossum umbratile
Geoglossum vleugelianum
Glutinoglossum glutinosum
Hemileucoglossum elongatum
Hemileucoglossum littorale
Trichoglossum hirsutum
Trichoglossum rasum
Trichoglossum tetrasporum
Trichoglossum walteri
Microglossum clavatum
Microglossum fuscorubens
Microglossum griseoviride
Microglossum nudipes
Microglossum nudipes aff.
Microglossum olivaceum s. str. HDI
Microglossum pratense
Microglossum rufescens
Microglossum tenebrosum
Microglossum truncatum

Other grassland fungi (*Camarophyllopsis*, ***Dermoloma***, ***Hodophilus***, *Pseudotricholoma*)
Camarophyllopsis schulzeri
Dermoloma cuneifolium
Dermoloma josserandii
Dermoloma magicum
Dermoloma pseudocuneifolium
Hodophilus atropunctus
Pseudotricholoma metapodium

bold = recorded in Surrey HDI = High Diversity Indicator

Annex 3. Waiting List of potential, as yet unverified sites of future interest

site name:	hectad:	GR (centroid):	provisional qualifying UK Hab 2.0 habitat:	existing site designations:	owner/occupier (if known):
Elmbridge					
Claygate, Wingham Court grounds	TQ16	TQ154641	g3c?		private
Cobham, R. Mole floodplain fields	TQ16	TQ100604	g3c/fg-m?		private
Esher, Arbrook House grounds	TQ16	TQ13996339	g3c?		private
Fairmile golf range	TQ16	TQ114615	g1a/d?		private
Weybridge, Wey floodplain W of railway	TQ06	TQ062634	g3c?		private
Guildford					
Ash, cemetery	SU85	SU897509	g1a/d?		CoE
Ash, Ash Centre playing field	SU85	SU895515	g1a?		LA
Ash, Shawfield Lane Meadows (E)	SU85	SU886504	g3c?		BVCP
East Horsley, Wellington Meadow	TQ05	TQ095527	g3c?		PC
Holmbury St. Mary, Hurtwood cricket pitch	TQ04	TQ108438	g1a/d?		PC?
Mytchett, Snakey Lane Wildlife Area	SU85	SU887540	g3c?		BVCP
Normandy Community Orchard	SU95	SU921521	g3c?		PC
Seale, grassland E of Seale Lodge sandpit	SU85	SU89454782	g1a?		private
Worplesdon, Blanket Mill flood-meadow	SU95	SU967546	g3c/fg-m?		private
Worplesdon, field at Rye Farm (Ellis's Field)	SU95	SU976546	g3c?	SNCI	private
Worplesdon, Marne's Field	SU95	SU975547	g3c?	SNCI	private
Mole Valley					
Ashtead, Larchfield Meadow	TQ15	TQ196573	g2c?		private
Betchworth House estate & park	TQ24	TQ209490	g3c/fg-m?		private
Betchworth, Wonham Park	TQ24	TQ219496	g3c damp?		private
Hookwood, Gatwick Park hospital grounds	TQ24	TQ271424	g3c?		NHS
Horley, R. Mole floodplain S of Westvale Rd	TQ24	TQ266442	g3c/fg-m?		private
Leatherhead, roughs W of M25	TQ15	TQ176566	g2c?		SCC
Little Bookham, field W of Maddox Lane	TQ15	TQ12075614	g3c?		private
Ockley, fields at New House Farm	TQ13	TQ140389	g3c?		private
Mole Valley-Reigate & Banstead					
River Mole floodplain, Ricebridge-Sidlow	TQ24	TQ225486-TQ257470	g3c/fg-m?		private
Reigate & Banstead					
Banstead, Ruffett Meadow	TQ25	TQ254586	g2a/c?		private
Salfords, field N of Honeycrock Lane	TQ24	TQ282468	g3c?		private
South Park, field S of Slipshatch Rd.	TQ24	TQ241482	g3c?		private
Woodmansterne, grasslands between Solom's & Scratch Woods	TQ25	TQ271590	g2a/c?		private
Runnymede					
Addlestone, fields E of M25 J11	TQ06	TQ040650	g3c/fg-m?		SCC/NH?
St. Ann's Heath, wet pastures	TQ06	TQ006679	g3c/fg-m?		private
Woodham, Holme Farm	TQ06	TQ033625	g3c?		private
Spelthorne					
Shepperton, Renfree Way roughs	TQ06	TQ072671	g3c?		private
Surrey Heath					
Chobham, field E of Dunstall Green	SU96	SU993627	g3c damp?		gas utility?
Chobham, Wentworth Place	SU96	SU964636	g1a/d?	SNCI	private
Chobham, fields E of Chobham Park Farm	SU96	SU988624 SU990626	g3c damp?	SNCI?	private
Tandridge					
Blindley Heath, fields to SE of Pond Farm	TQ34	TQ373445 TQ370440	g3c?		private
Caterham, Barber's Meadow	TQ35	TQ338541	g2a/c?		private
Caterham, fields at Beech Farmland E & W	TQ35	TQ33085377 TQ329538	g2a/c?		private
Caterham, grassland at Birch Wood	TQ35	TQ351562	g2a/c?		private
Caterham, Tillingdown Hill & valley	TQ35	TQ345564 TQ351558	g2a/c?		private
Caterham, White Knobs	TQ35	TQ343547	g2a/c?		private

Copthorne, field NE of Effingham Road	TQ34	TQ32634085			private
Dormansland, field N of Dencher Wood	TQ44	TQ406447	g3c?		private
Dormansland, fields E of Dormans station	TQ34	TQ397415	g3c?		private
Farleigh, fields S of Little Farleigh Green	TQ35	TQ379599	g3c?		private
Haxted, Eden Brook floodplain W of Dwelly Lane (inc. Mill Stream flood-meadow)	TQ44	TQ416458	g3c/fg-m?		private
Hurst Green, field E of The Diamond PH	TQ45	TQ405502	g3c?		private
Hurst Green, field to E of Holland Rd.	TQ44	TQ404499	g3c?	SNCI	private
Limpsfield, Moorhouse Bank, old sports field	TQ45	TQ430530	g1a/d?		private
Lingfield Cernes hinterland (non-SSSI)	TQ44	TQ413448 TQ410449	g3c/fg-m?		private
Lingfield, fields to S of Bowerland Farm	TQ34	TQ396453	g3c?		private
Oxted chalk-pit	TQ35	TQ382543	g2a/c?		private
White Post Common	TQ35	TQ332501	g3c?		PC?
Whyteleafe Rec.	TQ35	TQ341587	g2a/c?		SCC?
Waverley					
Bowlhead Green, fields nr. Begley Farm	SU93	SU903371	g3c?		private
Chiddingfold, Pheaton's Copse fields, Pook Hill	SU93	SU946350	g3a/c?		private
Chiddingfold, Ramster Gardens arboretum	SU93	SU948333	g3c*		private
Cranleigh, field E of Knowle Lane	TQ03	TQ057381	g3c?		private
Cranleigh, marshy fields S of Book Hurst	TQ03	TQ074389	g3c damp?		private
Dunsfold, field W of Knightons Lane	TQ03	TQ011343	g3c?		private
Elstead, Westbrook House fields	SU84	SU891429	g1a/d?		private
Ewhurst, Parliament Farm fields	TQ03	TQ095389	g3a/c?		private
Farnham, West Street cemetery	SU84	SU832463	g1a/d?		CoE
Hale, Farnham Park cemetery	SU84	SU847483	g1a/d?		private
Hale, St. John's churchyard & environs	SU84	SU849483	g1a/d?		CoE
Haslemere, field at Invall	SU93	SU900355	g1a?		private
Haslemere, fields E of Bunch Lane	SU83	SU898337	g1d/g3c?		private
Haslemere, meadow S of Keffolds Farm	SU93	SU908346	g1a?		private
Haslemere, Swan Barn Farm	SU93	SU907329	g3c?		NT
Milford, Oxenford Grange watermeadows	SU94	SU937432	g3c/fg-m?		private
Thundry Meadows, Badgers Bank	SU84	SU896442	g3c?	SSSI?	private
Wonersh, field N of Barnett Hill House	TQ04	TQ025461	g1a/d?		private

Surrey Nature Partnership
Healthy Environment | Healthy People | Healthy Economy

SURREY IMPORTANT GRASSLANDS INVENTORY 1.0

Surrey Nature Partnership

Reference: **Waite**, **M J**. (ed.). 2026. *Surrey Important Grassland Inventory* 1.0.
Surrey Nature Partnership